

Chiasson, Frédéric, Caroline Traube, Clément Lagarrigue, and Stephen McAdams, Koechlin's Volume: Perception of Sound Extensity among Instrument Timbres from Different Families, *Musicae Scientiae* (volume: 21, issue: 1) pp. 113–31. Copyright © the authors 2016. DOI: [10.1177/1029864916649638](https://doi.org/10.1177/1029864916649638).

Koechlin's volume: Perception of sound extensity among instrument timbres from different families

Frédéric Chiasson, Caroline Traube, Clément Lagarigue
Faculté de musique, Université de Montréal
Montréal, Québec, Canada

Stephen McAdams
Schulich School of Music, McGill University
Montréal, Québec, Canada

Musicae Scientiae
(in press)

Abstract

Charles Koechlin's *Traité de l'orchestration* ascribes different dimensions to timbre than those usually discussed in multidimensional scaling studies: “volume” or *grosseur*, related to extensity (the sound's perceived size), and intensity, related to loudness. Koechlin also provides volume rankings for orchestral instruments in different registers. Studies show that humans, as well as several animal species, perceive extensity for many sound sources, but none has demonstrated its relevance for musical instruments from different families. To test extensity, samples of seven orchestral instruments equalized in pitch, but not in loudness, were used. Task 1 required participants to order eight sets of samples on a largeness (*grosseur*) scale from "less large" (*moins gros*) to "larger" (*plus gros*). Task 2 required them to quantify the sounds' largeness compared to a reference sample on a ratio scale. Both studies show that participants share a common extensity perception for instrument timbres of different families that is very similar to Koechlin's proposed scale. This perception seems to be related to spectral shape and particularly to acoustic energy in the lower frequencies. Perception of this attribute is unrelated to musical training, native language, and the presence of minor hearing loss, which suggests that extensity could be a universal attribute of timbre perception that is useful in orchestration practice and theory.

Keywords: volume, extensity, largeness, auditory size, timbre, orchestration, Koechlin

Koechlin's *Volume*: Perception of Sound Extensivity Among Instrument Timbres from Different Families

Many researchers, such as Hajda, Kendall, Carterette and Harshberger (1997), have pointed out that the official ANSI psychophysical definition of timbre—“that attribute of auditory sensation in terms of which a subject can judge that two sounds similarly presented and having the same loudness and pitch are dissimilar”—is problematic for its study. Still, most multidimensional scaling (MDS) studies on timbre have based their research methodology on this definition, equalizing loudness and pitch to find the acoustic factors that contribute to timbre perception (e.g., McAdams, Winsberg, Donnadieu, De Soete & Krimphoff, 1995). Hadja *et al.* (1997) reviewed many MDS studies of timbre and determined that spectral centroid was the most salient acoustic measure for timbre perception in continuous sounds. Peeters, Giordano, Susini, Misdariis and McAdams (2011) conducted a correlational analysis, followed by hierarchical clustering, on a large set of acoustic descriptors of timbre across pitch, dynamics and articulation on a database of about 6000 musical instrument sounds and identified ten classes of primary descriptors. The four largest groups included: 1) the central tendency of time-varying spectral properties (e.g., spectral centroid), 2) the temporal variability of the time-varying spectral properties (e.g., spectral variation), 3) global energetic properties (e.g., energy of harmonic and noise components) and properties of the temporal envelope (e.g., attack and decay times), and 4) descriptors of signal periodicity (e.g., noisiness and inharmonicity).

Obviously, musicians do not use such acoustic terminology to describe the perceptual attributes of instrumental sounds but rather use a more intuitive set of verbal descriptors referring to the roundness, sweetness, sharpness, brightness, openness or largeness of sound. The French composer Charles Koechlin, author of an important orchestration treatise (1954-1959), has proposed a unique orchestration system that uses general timbre descriptors to explain the sound of diverse orchestrations by Bach, Mozart, Beethoven, Berlioz, Wagner, Bizet, Debussy, Ravel, Stravinsky, Milhaud, and Schoenberg. One crucial timbre dimension Koechlin discusses in this

treaty is designated by the term *volume* (Koechlin, 1954-1959, vol. I, p. 220). This is the perceptual attribute that we are investigating in this study.

Koechlin's orchestration System

Koechlin uses *volume* and *intensité* as the basic timbre descriptors for a complex and highly refined orchestration “system” (Koechlin, 1954-1959, vol. I, p. 220). He does not describe this system as such, but a careful reading of his *Traité de l'orchestration* shows coherent relations among the timbre descriptors used in a way that makes it possible to reconstitute the system. Figure 1 shows the relations among descriptors for orchestration, according to the musical “object” defined: instrument, orchestral layer or the whole orchestra (Chiasson, 2010).

[Insert Figure 1 here]

Volume corresponds to the extensivity of an instrument sound, the space it seems to occupy in the auditory scene. Intensity (*intensité*) corresponds to its perceived loudness or strength. Both qualities are used to evaluate the instrument timbre's transparency (*transparence*) or its opposite, density (*densité*) (Koechlin, 1954-1959, vol. I, p. 220). A timbre is dense when it has a high intensity with a small volume, whereas it is transparent when it has a low intensity and a large volume. When other musical parameters are similar, a dense timbre is heard in the foreground and a transparent timbre in the background, so these parameters may affect figure/ground perception in orchestral music.

Timbres with similar volume and intensity are balanced (*équilibré*) (Koechlin, 1954-1959, vol. I, p. 223). This balance is required for instruments to blend (*fondre*) into an orchestral layer (*plan orchestral*). A high degree of blend (*fondue*) between similar instruments (most of the time between instruments from the same family, like strings) playing in similar registers (*tessitures similaires*) creates a homogeneity (*homogénéité*) of the layer (Koechlin, 1954-1959, vol. III, p. 3).

A high level of blend and homogeneity in the layers of the orchestration and a strong use of instruments with large volume create a perception of “fullness” (*plénitude*) for the whole

orchestration. Koechlin calls this “material fullness” (*plénitude matérielle*). On the other hand, a fullness of the compositional writing (*plénitude d'écriture*) is heard when the orchestration uses voice doublings (*doublures*) at different intervals, counterpoint in an orchestral layer and instruments playing in the register called “submedium” (*sous-médium*) by Koechlin, which is the octave below middle C. Finally, brightness or vividness (*éclat*) is heard when, on the contrary, some orchestral layers are made of single instruments without doublings, which Koechlin calls “pure timbres” (*timbres purs*) (Koechlin, 1954-1959, vol. III, pp. 7-10).

Koechlin's volume scale and the scientific literature

Koechlin provides a scale of the relative volumes of traditional orchestral instruments, as shown in Table 1, while stressing the fact that volume can change with register, dynamics, the number of musicians playing the same part, and even the music played (Koechlin, 1954-1959, vol. I, p. 288).

[Insert Table 1 here]

If Koechlin's *intensité* or loudness is intuitively obvious for both music psychologists and musicians, volume does not show such consensus. Although volume is considered to be a common musical fact by French authors (Koechlin, 1954-1959; Schaeffer, 1966) and German authors, under the German term of *Tongröße* (“tone size”) (Stumpf, 1890; Albersheim, 1939), most timbre studies in the field of psychoacoustics do not mention any volume- or acoustic-size-related dimension (Hajda *et al.*, 1997; Giordano & McAdams, 2006). However, one MDS study on violin timbre as perceived by musical experts does have a “narrow/full” dimension that indirectly relates to the “thin” and “large” descriptors of the volume scale (Štěpánek, 2006). Also, Kendall & Carterette (1993) have listed verbal descriptors that seem to relate to volume in Walter Piston's *Orchestration*—“voluminous” (violin), “full” (cello, trombone, heckelphone), “broad” (horn), etc. These descriptors, found in most orchestration treatises (e.g., Adler, 2002; Blatter, 1997; Piston, 1955; Rimsky-Korsakoff & Steinberg, 1914), are mostly specific to an instrument or a unique combination of instruments. Verbal descriptors related to volume—full,

open, thick versus small, closed and thin—are used by performers in pedagogical or rehearsal contexts (Traube & Depalle, 2004).

It should be noted that the concept of volume, defined as “an apparent largeness and extensiveness,” has been investigated by the pioneer of psychoacoustics S. S. Stevens. With different collaborators, he conducted a first group of studies on the volume perception of pure sounds. They observed that the volume of pure sounds increased when loudness increased and pitch decreased (Stevens & Davis, 1938; Terrace & Stevens, 1962), and that loudness was a product of volume and density (Stevens *et al.*, 1965), the same relation proposed by Koechlin (1954-1959).¹

In this paper, we have chosen to use the term *extensity* to refer to the concept of volume as defined by Koechlin and as used by Stevens in his studies. Stevens himself acknowledges the current ambiguity of the word, because when “the radio-engineer speaks of [...] the ‘volume control’ of a radio set, [...] he means [...] what we should probably call intensity” (Terrace & Stevens, 1962, p. 597).

The term *extensity* can be found in early texts in the psychology of perception. For example, Edward Titchener in 1896 defined extensity as an “impression that a sensation or image is more or less spread out in space” (cited in Hergenhahn & Henley, 2013, p. 261). Erickson (1975, p. 144) uses the term extensity as the “bigness“, “spaciousness, broadness [and] diffuseness of a sound”. In the field of auditory perception, the expression *auditory size* has also been used, such as in recent studies using human voice stimuli (Smith, Patterson, Turner, Kawahara & Irino, 2005), noise bands (Cabrera & Tilley, 2003), and musical instruments from the same family (Dinther & Patterson, 2006). Auditory size perception has also been investigated in other animal species (Fitch, 1997; Huron, Kinney & Precoda, 2006; Reby & McComb, 2003). We have adopted the term *extensity* as it is less ambiguous than volume and it refers to a sensation that is inherent to the sound and independent of the size of the sound-producing object itself.

General Methods

The purpose of our first experiment was to verify whether participants' rankings of extensity differences among sounds of musical instruments from different families on an ordinal scale correspond to the relations theorized by Koechlin (Exp. 1). We also wanted to know whether participants can quantify extensity in instrument sounds using a ratio estimation procedure (Exp. 2) as has been done with pure sounds in Terrace and Stevens (1962).

Stimuli

Koechlin developed his volume scale based on his own experience hearing musical instruments during numerous concerts. The major issue concerning the instrument samples was to find a sample bank that kept the loudness differences between instruments as they would be heard in a concert hall. The oboe, for instance, rarely plays louder than a trumpet in a concert. However, this loudness ratio is often lost with the loudness equalization often required in MDS studies. Due to the different recording conditions used for each instrument to produce instrumental sound databases such as the McGill University Master Samples (Opolko & Wapnick, 1989), the oboe sounds louder than the trumpet, which is unrealistic for an orchestrator. The difference in loudness between concert and laboratory situations would probably create a significant difference in the extensity of each instrument.

In the Music Instrument Samples database from the Electronic Music Studio of the University of Iowa (Fritts, n.d.), each instrument is recorded at various dynamic levels as controlled by the player, from *pianissimo* to *fortissimo*. The sound pressure level of the samples is not normalized, and this is why we chose to work with this database.

All instruments were recorded in the Anechoic Chamber in the Wendell Johnson Speech and Hearing Center at The University of Iowa using a Neumann KM 84 cardioid condensor microphone (Georg Neumann GmbH, Berlin, Germany) and a Mackie 1402-VLZ mixer (Loud

Technologies, Inc., Woodinville, WA) with a Panasonic SV-3800 DAT recorder (Panasonic Corp., Osaka, Japan). All instruments were played at a distance of 1.5 m from the microphone.

Seven instruments from among those covered in Koechlin's volume scale were used: violin, flute, oboe, B \flat clarinet, alto saxophone, C trumpet and French horn (hereafter simply “horn”). We used samples of these instruments playing F or B in octaves 3, 4 or 5 ($A_4 = 440$ Hz), according to the available range of each instrument (Table 2). The horn playing F3 is absent, because the horn samples in octave 3 were not available. Each instrument was recorded playing at a dynamic marking of *mf*, the choice of loudness being left to the player and following the instrument's capacities. All instruments were played without vibrato and were recorded in mono, 16 bit, 44.1 kHz, AIFF format.

[Insert Table 2 here]

All samples were one second in duration for this study, with a linear fade-out over the last 100 ms. Onsets were kept in their original form for all samples. No loudness equalization was performed to keep the natural loudness differences among the instruments and to be closer to the “natural” concert situation. Pitch equalization was used, however. The mean pitch deviation from equal temperament was calculated for each sample with Praat (University of Amsterdam, Amsterdam) and then transposed to equal temperament using the Elastic Pitch tool in Pro Tools 8 LE (Bias, Inc., Petaluma, CA). The mean correction was 13.2 cents. Apart from the horn B3 sample, tuned down by 116 cents because it was produced with the C4 sample, the maximum correction was 32.1 cents.

Apparatus

Participants were seated in a sound-attenuated room. The samples were played on two loudspeakers separated by 155 cm, facing the participant, whose chair was approximately 75 cm from the line between the two speakers. The middle of each speaker was 112 cm high, approximately at the same level as the participant's head. The computer screen was at the exact middle of the line between the speakers. The trumpet B5 sample, which was the loudest, was calibrated to 75 dB SPL (C-weighting) for all participants using a Quest SE SoundPro DL 1/3

Octave level meter (3M Detection Solutions, Oconomowoc, WI) on the slow rate setting, positioned at the same position as the center of the listener's head when seated.

Participants

The 80 participants consisted of 40 native francophones and 40 native anglophones, with equal numbers of musicians and nonmusicians in each group. Among the anglophones, 22 were not fluent in French in terms of listening, speaking, reading or writing. Forty participants were professional musicians or music students with at least an undergraduate level, and 40 were either amateur musicians not studying in music or nonmusicians. Fifty-eight participants had normal hearing with no hearing loss of more than 20 dB SPL in both ears as measured with a standard audiometric procedure (ISO 389-8, 2004; Martin & Champlin, 2000), and 22 had some hearing loss over 20 dB or other minor hearing problems (mostly tinnitus). Before doing both experiments, all participants signed an informed-consent form and filled out a questionnaire concerning language, hearing issues, music listening habits, instrumental practice, and music theory knowledge. After the experiments, participants were questioned by the experimenter concerning their listening strategies and strategies for performing the tasks in both experiments. All participants took part in both experiments.

Experiment 1

Method

In the first experiment, participants were asked to rank eight sets of samples in order of *gros* (for francophones) or “largeness” (for anglophones) from *moins gros* or “less large” to *plus gros* or “larger”. As mentioned earlier, the word “volume” is now confused with acoustic intensity or loudness (which we observed during the pilot study), so we replaced Koechlin's term “volume” by *gros* in French, and “largeness” in English. The instructions were presented on the software interface, programmed in Max/MSP (Cycling 74, San Francisco, CA).

For each series, the experimental interface presented a number of boxes at the top of the screen and an equivalent number of baskets below. Each box represented a sample to be ranked in this series and the baskets represented the largeness ranks. The initial order of sample boxes

was randomized for each series. Participants had to double-click on the box to hear the sample and to drag the box to a basket. They were asked to put all boxes in order of largeness in the baskets, one box per basket. They were also asked to listen to all samples during the process and to confirm their volume ranking by listening to all samples in order before proceeding to the next series.

Before the main set, a practice set was performed with the experimenter present, with samples of horn, flute, oboe, saxophone and violin playing B4. The experimenter gave no feedback concerning the accuracy of answers according to Koechlin's scale. Then in the main set, participants ranked six single-pitch sets, then two trans-pitch sets (Table 2). The presentation orders of the six single-pitch sets and of the two trans-pitch sets were randomized for each participant.

Results

The median rankings for each instrument in each sound set are shown in the box-and-whisker plots in Figure 2. There are too few points to perform reliable correlations between the median rankings and the ranks along Koechlin's volume scale. However, as an indication, rank correlations are significant ($r^2 \geq .92$, $p < .01$) for B4, F5, B5 and marginally so ($r^2 = .76$, $p = .078$) for TransB. The relations are less clear for F3, B3, F4 and TransA ($r^2 < .63$, $p > .10$).

[Insert Figure 2 here]

We used the ranks for mixed ANOVAs on each series with Instrument as repeated measure and native Language (English, French), Musicianship (musician, nonmusician) and Hearing (no hearing problems, hearing problems) as between-subjects factors. Mauchly's test of sphericity was significant at the .001 level for all eight analyses, so multivariate statistics were used, which do not assume sphericity. The Instrument main effect was highly significant for all eight analyses, Wilks' $\lambda < .325$, $p < .001$, $\eta_p^2 > .675$. The only interaction with Instrument that was significant was the Instrument \times Language \times Musicianship \times Hearing interaction for the F3 series,

Wilks' $\lambda=.835$, $F(2,71)=7.0$, $p=.002$, $\eta_p^2>.165$. None of the between-subjects main effects or their interactions were significant. However, Shapiro-Wilk tests showed that none of the resulting distributions—according to language, musicianship or hearing—were normal, because of the rank data. Levene tests also showed that only a few series had the necessary homogeneity of variance to respect the assumptions underlying ANOVA. For this reason, we also performed Friedman and Mann-Whitney non-parametric tests on all data from Experiment 1. Friedman's test was used to determine if the ranking distributions were the same for each instrument. The tests showed that all series had different distributions for each instrument ($p < 0.001$ in all sets), which is consistent with ANOVA results. A Mann-Whitney test was used to determine if the mean rankings for each instrument differed across levels of the between-subjects factors Language, Musicianship and Hearing. None of these comparisons between language groups for each instrument in each series were significant after correction for multiple tests using the Bonferroni-Holm method.

In sum, the rankings given by the participants globally varied across instruments similarly to Koechlin's scale for the mid to high registers, with a few lower-than-expected values for the flute and horn in some series. The results were not reliably affected by Language, Musicianship or Hearing. This would suggest that extensity perception among instruments is not something that is learned directly by music education or indirectly by a larger cultural education related to a language or an historical period (Koechlin wrote his treatise during the 1930s even though it wasn't published until the 1950s). We hypothesize that it likely involves a lower-level timbre recognition process that is part of a more general sound source perception process. To examine further the extent to which and the precision with which listeners perceive extensity, we designed another experiment with the same sounds using psychophysical scaling, which aims to quantify more precisely the perceived degree of extensity.

Experiment 2

Method

On each trial, participants heard a reference sound followed by a comparison sound and were asked to give a numerical rating on a ratio scale of largeness or *grosseur* of the comparison sound with respect to the reference sound. The reference sound was the sample of clarinet playing B4, because it was in the middle of Koechlin's scale and one of the pitches in the middle of our pitch range, in a series in which the instruments had the greatest diversity of extensivities according to Koechlin's scale (see Figure 1). It was assigned a largeness/*grosseur* value of 100. Participants were not informed about the nature of the reference sound. Each of the 40 sounds from Experiment 1 was presented as a comparison sound, including the reference sound. For instance, if the test sound seems twice as large as the reference sound, write 200; if it sounds one third of the reference sound largeness, write 33. Participants were asked to evaluate only the sound largeness, even if there were different loudnesses and pitches in some series.

The instructions were presented via a Max/MSP interface. A practice set of three samples (horn at B4, flute at B3 and saxophone at F5) presented in randomized order was rated in the presence of the experimenter, during which no feedback about the correctness of the answers was given. During the main set, participants rated each of the 40 samples from Experiment 1 once.

Results

Similarly to Experiment 1, the median results in Experiment 2 globally follow Koechlin's proposed volume scale, still with some exceptions for flute and horn as in Experiment 1 (Figure 3). Again, except for the F3 and TransA sets, almost all median values correlate well with Koechlin's volume scale. To have a more accurate estimation of the correlation, we used the median values from Experiment 2 to calculate the instrument ranks. We then computed Spearman correlations between the median ranks of each series and Koechlin's scale. The correlations between predicted extensity scales according to Koechlin and participants' mean largeness scales are significant for all groups and explain 73% to 83% of the variance in the data.

In line with S. S. Steven's (1971) power law, we used the log of each ratio estimation as the dependent variable for mixed ANOVAs on each sound set, with Instrument as repeated measure and native Language (English, French), Musicianship (musician, nonmusician) and Hearing (no hearing problems, hearing problems) as between-subjects factors. Mauchly's test of sphericity was significant at the .001 level for all eight analyses, so multivariate statistics were used, which do not assume sphericity. This time, Shapiro-Wilk tests showed that most of the data were normally distributed for levels of the factors Language, Musicianship and Hearing. Levene tests also showed that most of the data respected the homogeneity of variance assumption. Levene tests that rejected the null hypothesis of homogeneity of variance were on Instruments related to Musicianship (clarinet in series F3, B3 and B5; flute in series F4, B4, F5 and B5; saxophone in series B4 and F5; horn in series F5; oboe in series B5). The main effect of Instrument was highly significant for all eight series, Wilks $\lambda < .443$, $p < .001$, $\eta_p^2 > .557$. Instrument \times Language interactions were significant only for series B4, $\lambda = .83$, $F(6,67) = 2.30$, $p = .045$, $\eta_p^2 = .170$, and F5, $\lambda = .781$, $F(6,67) = 3.13$, $p = .009$, $\eta_p^2 = .219$. The Instrument \times Musicianship \times Hearing interaction was significant only for series B3, $\lambda = .78$, $F(6,67) = 3.09$, $p = .01$, $\eta_p^2 = .217$, and TransA, $\lambda = .76$, $F(5,68) = 4.30$, $p = .002$, $\eta_p^2 = .240$. The effect of these variables is fairly small compared to that of the Instrument main effect, and they don't seem to be systematic enough in pattern to interpret. T-tests comparing groups for Musicianship and Hearing on each instrument in each series revealed only one significant comparison after Bonferroni-Holm correction: the violin in the B4 series for the two Language groups, $t(78) = -2.39$, $p < .001$.

The ANOVAs did not reveal any significant influence of Musicianship and Hearing factors on the data ($p > .19$ in all cases). Mann-Whitney tests revealed an influence of Musicianship only for saxophone in the F3 series ($p = .004$), as well as a significant influence of Hearing on trumpet extensivity in five of the eight series ($p < .047$ in all five cases) and on flute ($p = .043$). In all concerned series, trumpet is heard as smaller by participants with hearing problems than by participants with normal hearing.

Again, Instrument is the most salient factor in the results for Experiment 2. Although Language has a significant effect in some cases, it does not have a great effect on the extensity ratio scaling of instruments. These results are consistent with the results of Experiment 1.

Discussion

As for Experiment 1, the results from Experiment 2 show that except for the F3 and TransA series, participants have a common relative extensity perception of the instrument samples, which is again correlated with Koechlin's scale (Table 4), and the rank correlation again increases with pitch. Koechlin's scale seems to be able to roughly predict participants' extensity rankings, even when they are asked to give numerical estimations of extensity to instruments in random order, with no grouping according to series.

Instrument is the most significant factor for extensity rankings. While most instrument rankings follow Koechlin's scale, there is one strong discrepancy between the participants' scale and Koechlin's: the place of the flute in its low register. Koechlin insists in his *Traité* that the flute in its low register is as large as the trumpet at the same pitch. However, participants estimated the flute's extensity as much lower than most instruments, even as the lowest for the B3 series. This difference is observed for all participant groups.

A slighter discrepancy can also be seen between participants' and Koechlin's scales for the place of the horn. According to Koechlin, the horn should be the instrument sounding the largest among our set. In our experiments, it is still at the largest end of the scale, but shares its position with trumpet and saxophone. Thus, the horn sounds somewhat smaller for our participants than for Koechlin.

With the actual data and settings, the reason for the discrepancies in extensity rankings for flute and horn is hard to determine. It could be caused by a difference in the perception of these instruments between the experimental and normal concert contexts. The places of both musicians and listeners in the concert hall and the hall's own acoustic characteristics affect the perceived timbre and could enlarge the extensity of these instruments. It could also be caused by

Koechlin's particular way of perceiving flute and horn extensivities, which are not shared by other listeners.

Language is also a significant factor for most series, but this influence is small. In Experiment 2, Hearing was a significant factor affecting Trumpet largeness ratings, but again, this influence is small. These results provide evidence for a common extensity perception by Koechlin and our listeners that is only minimally affected by language, musical training or minor hearing problems. The relation between median extensity scalings and Koechlin's volume rankings show a very similar pattern to those from Experiment 1 in spite of the differences in the tasks (ordinal ranking and ratio scaling). Indeed the Spearman rank correlation coefficients between the median values in the two experiments vary from .87 to 1.00 across the eight series. Table 3 juxtaposes the rankings derived from Koechlin's *Traité* and from Experiments 1 and 2. Values from the experiments that are not statistically different are shown in the same rank categories.

[Insert Table 3 here]

Acoustic correlates of extensity

Having demonstrated differences in extensity across instruments and pitch registers, we now turn our attention to establishing an acoustic basis for the perception of extensity. A Principal Component Analysis (PCA) with Varimax rotation and Kaiser normalization was used first to determine the number of independent latent variables representing a set of 12 acoustic descriptors. We then compared them to the median results from Experiment 2, because the numerical nature of extensity ratio scaling gives a more precise correlation with the acoustic descriptors used, and at any rate, the median results in both experiments are highly correlated. In concordance with the literature on psychophysical scaling (Stevens, 1975), we used the logarithm (base 10) of the extensity values. The descriptors included loudness (in sones), spectral centroid (a measure of the amplitude-weighted average frequency of the spectrum, hereafter referred to as absolute spectral centroid), fundamental frequency (F_0), the spectral centroid/ F_0 ratio (centroid expressed in harmonic rank independently of the fundamental frequency, referred hereafter as

relative spectral centroid), spectral flux (a measure of spectral variation over time), spectral rolloff (frequency below which 95% of a sound's frequency content is located), spectral spread, spectral skewness, spectral kurtosis, attack time, even/odd harmonics amplitude ratio and irregularity across the amplitudes of the frequency components (spectral irregularity). All of these descriptors were computed using the MATIMBRE Matlab toolbox (Barthet, 2008).

The PCA analysis rejects components that have eigenvalues less than 1. The analysis converged in four iterations and extracted three components that explain 39%, 20% and 13% of the variance, respectively, for a total variance explained of 72%. The loadings of each descriptor on each component are shown in Table 4. The descriptors that load the strongest on principal component 1 (PC₁) include spectral kurtosis, spectral spread, and spectral skewness, with weaker contributions from absolute spectral centroid and loudness. Note that the value of this latent variable increases with increases in absolute spectral spread and spectral centroid and with decreases in spectral kurtosis, spectral skew and loudness. PC₂ represents spectral roll-off and spectral irregularity (decreasing), and PC₃ includes F₀ and relative spectral centroid (decreasing), with weaker contributions by attack time (decreasing), spectral flux, and even/odd ratio (decreasing).

[Insert Table 4 here]

Entering the three PCs as independent variables into a multiple regression onto the log median extensity values (E) of all instrument samples at all pitches from Experiment 2 yields the following equation:

$$\log(E) = 2.08 - .74 PC_1 - .32 PC_2 + .06 PC_3. \quad (1)$$

This equation suggests that the acoustic descriptors that increase perceived extensity are those represented by PC₁ and PC₂. Extensity increases most strongly with increases in spectral kurtosis, spectral skewness towards low frequencies, and to a lesser extent in loudness and spectral irregularity, and it increases with decreases in spectral spread, spectral centroid and spectral roll-off. Acoustic descriptors that decrease largeness perception are spectral centroid and

spectral spread. Note that the inverse relation between spectral centroid and largeness perception was also observed by Stevens *et al.* (1965).

With the acoustic factors extracted from the PCA, we modeled two idealized spectral envelopes that relate to a large sound and a small sound. The spectral envelope of a large sound has a sharp peak of high amplitude and low frequency skewed toward the higher frequencies, with energy concentrated around that peak. The spectral envelope of a small sound has a fairly symmetrical peak of low amplitude and higher frequency, with energy being spread more broadly over the spectrum (Figure 4). When put in order of perceived largeness, graphs of the spectra of instrument samples in the F4 series show a transition of spectral envelopes from small sound to large sound envelope models, suggesting the relevance of these model curves (Figure 5). The concordance of these models with the spectral envelopes of our samples suggests that extensity might be a function of the energy of lower absolute frequencies.

To verify this hypothesis further, we performed a PCA with the specific loudnesses extracted from auditory filters equally spaced according to ERB-rate (equivalent rectangular bandwidth scale) for each instrument sample (Moore & Glasberg, 1983). Two significant PCs were extracted. PC_a was related positively with higher frequency bands (>583 Hz) and slightly negatively with lower frequency bands (<583 Hz). PC_b was related positively with lower frequency bands (<897 Hz) and slightly negatively with higher frequency bands (>897 Hz).

We then ran a stepwise multiple linear regression on both PCs and the log median extensity values from Experiment 2. The resulting equation is:

$$\log(E) = 2.08 + .094 PC_a + .11 PC_b . \quad (2)$$

This equation shows that both higher (PC_a) and lower frequencies (PC_b) contribute to the extensity value, but that lower frequencies (PC_b) have a slightly higher contribution to the extensity value. This equation suggests that our model envelopes go in the right direction, although they should be corrected to reflect the influence of higher frequencies and overall signal amplitude.

However, because of the irregular nature of the samples, it is difficult to quantify more precisely how the specific loudness in each frequency band contributes to the perceived extensity of each sample. Further research with synthesized sounds that vary according to the significant acoustic descriptors found in these experiments would be needed to confirm this interpretation.

The methodology used in this study demonstrates that participants can hear differences of largeness consistently even when samples have loudness differences. Preserving loudness differences should be interesting in some experimental situations like ours, where it is important to keep timbres as close as possible to a concert situation. As we have seen, to equalize the samples' loudnesses would have meant changing the spectra of the samples, thus altering the differences in extensity among them.

These results also suggest that it is not essential to have samples with equal pitches and loudnesses to have valid experimental methodology to test timbre perception. The present study shows that differences of a particular feature of timbre can be heard and discriminated even with differences of loudness when the participants are directed to do so, as postulated by Hajda *et al.* (1997). Another study by Marozeau, de Cheveigné, McAdams and Winsberg (2003) on the influence of pitch on MDS analyses of timbre, shows that differences in timbre can also be heard and discriminated among samples of different pitch, when the participants are asked to do so. Moreover, as our study suggests, differences in pitch and loudness *could* contribute to the perceived timbre difference due to their effects on acoustic parameters that affect timbre.

Finally, numerous experimental results among different animal species (Fitch, 1997; Huron et al., 2006; Reby & McComb, 2003) and with different sound sources (for experiments with humans; Cabrera & Tilley, 2003; Dinther & Patterson, 2006; Smith et al., 2005) tend to confirm, along with our own experiments, that extensity might be an auditory perception of sound that would exist beyond cultural learning and even human-specific audition. Extensity may be a hearing attribute that is useful for different biological behaviors such as judging the size of an unseen organism or sounding object and making decisions and acting appropriately based on the potential meaning of that perception. The adoption of this attribute for musical purposes

would thus be a deviation from its former ecological origins, extending it to a more general sense of sound extent with its implication in multiple sound sources through orchestration. As such, we now understand why this feature of hearing was so important for Koechlin (and many other classical musicians). A good understanding of extensity (or *volume*, as Koechlin called it) is crucial to give orchestrations “illusions” of “largeness”, “distance”, “depth” and of other evocative descriptions in Koechlin's *Traité de l'orchestration*. The art of orchestration becomes an art of creating plausible “illusions” of auditory scenes—analogueous to theatre and cinema creating “theatrical illusion”—by using limited auditory cues taken from real experience. For instance, a muted horn playing softly would be used to suggest a large musical call coming from afar, although the horn plays next to every other musician of the orchestra. Hence, using extensity wisely (among other attributes of hearing), a composer can create refined and detailed musical “scenes” in which many features can be heard by most listeners.

Conclusion

The purpose of this study was to verify the validity of Koechlin's mean *volume* (or largeness) scale in native French and English speakers with different musical backgrounds. The results show strong evidence that participants hear the relative extensity of instrument samples roughly as Koechlin has described, independently of native language, musical background or the presence of a small hearing loss in the high frequencies. High energy concentrated in low frequencies tends to increase perceived largeness, whereas low energy more spread out in higher frequencies tends to decrease it. Strong results consistent with other studies with different sound sources or animal species suggest that extensity is a part of a general collection of timbre perception processes, oriented to recognize properties of different sound sources. And composers have capitalized on its biological origins to serve musical purposes.

Funding

This work was supported by a grant from the Canadian Natural Sciences and Engineering Research Council (RGPIN 312774-10) and a Canada Research Chair awarded to Stephen McAdams. It was also supported by a grant from the Fonds de Recherche Québec—Science et

Culture awarded to Frédéric Chiasson and a grant from the Laboratoire Informatique, Acoustique et Musique (LIAM) to Clément Lagarrigue.

Notes

1. Stevens thought that density and brilliance might be a single acoustic quality (Stevens & Davis, 1938). Because brilliance is the perceptual correlate of spectral centroid (Hajda *et al.*, 1997) and most MDS studies use loudness-equalized samples (Hajda *et al.*, 1997; McAdams, 1993), volume may be indirectly related to the spectral centroid dimension if we consider the loudness equation from Stevens *et al.* (1965) according to which volume increases when density or brilliance decreases for sounds with equal loudness.

References

- Adler, S. (2002). *The Study of Orchestration* (3rd ed.). New York: W.W. Norton.
- Albersheim, G. (1939). *Zur Psychologie der Toneigenschaften [On the Psychology of Tone Quality]*. Straßburg: Heitz.
- Barthet, M. (2008). *De l'interprète à l'auditeur: Une analyse acoustique et perceptive du timbre musical [From performer to listener: An acoustical and perceptual analysis of musical timbre]*. PhD thesis. Aix, France: Université d'Aix-Marseille II. Retrieved from <https://tel.archives-ouvertes.fr/tel-00418296>.
- Blatter, A. (1997). *Instrumentation and Orchestration* (2nd ed.). Australie, Canada: Schirmer Books.
- Cabrera, D., & Tilley, S. (2003). Parameters for auditory display of height and size. *Proceedings of the 2003 International Conference on Auditory Display*, 29–32.
- Chiasson, F. (2010). L'universalité de la méthode de Koechlin [The universality of the method of Koechlin]. In P. Cathé, S. Douche & M. Duchesneau (Eds.), *Charles Koechlin : Compositeur et humaniste [Charles Koechlin: Composer and humanist]* (pp. 397-413). Paris: J. Vrin.

- Dinther, R. V., & Patterson, R. D. (2006). Perception of acoustic scale and size in musical instrument sounds. *Journal of Acoustical Society of America*, 120(4), 2158–2176.
- Erickson, Robert. (1975). *Sound Structure in Music*. Berkeley: University of California Press.
- Fitch, W. T. (1997). Vocal tract length and formant frequency dispersion correlate with body size in rhesus macaques. *Journal of Acoustical Society of America*, 102, 1213–1222.
- Fritts, L. (n.d.). *University of Iowa Musical Instrument Samples*. Mono recordings. Retrieved from <http://theremin.music.uiowa.edu/MIS.html> (pre-2012 version).
- Giordano, B. L. & McAdams, S. (2006). Material identification of real impact sounds: Effects of size variation in steel, glass, wood and plexiglas plates. *Journal of the Acoustical Society of America*, 119, 1171–1181.
- Hajda, J. M., Kendall, R. A., Carterette, E. C., & Harshberger, M. L. (1997). Methodological issues in timbre research. In I. Deliege & J. Sloboda (Eds.), *Perception and Cognition of Music* (pp. 253–306). Hove: Psychology Press.
- Hergenhahn, B. R., & Henley, T. (2013). *An Introduction to the History of Psychology*, 7th Edition. Boston: Wadsworth Publishing.
- Huron, D., Kinney, D., & Precoda, K. (2006). Influence of pitch height on the perception of submissiveness and threat in musical passages. *Empirical Musicology Review*, 1(3), 170–177.
- ISO 389-8. (2004). *Acoustics: Reference zero for the calibration of audiometric equipment—Part 8: Reference equivalent threshold sound pressure levels for pure tones and circumaural earphones* (Technical Report). Geneva, Switzerland: International Organization for Standardization.
- Kendall, R. A., & Carterette, E. C. (1993). Verbal attributes of simultaneous wind instrument timbres: II. Adjectives induced from Piston's *Orchestration*. *Music Perception*, 10, 469–502.
- Koechlin, C. (1954–1959). *Traité de l'orchestration, en 4 volumes [Treatise on orchestration, in 4 volumes]*. Paris: Max Eschig.

- Marozeau, J., de Cheveigné, A., McAdams, S. & Winsberg, S. (2003). The dependency of timbre on fundamental frequency. *Journal of the Acoustical Society of America*, 114, 2946-2957.
- Martin, F. N. & Champlin, C. A. (2000). Reconsidering the limits of normal hearing. *Journal of the American Academy of Audiology*, 11, 64-66.
- McAdams, S. (1993). Recognition of sound sources and events. In S. McAdams & E. Bigand (Eds.), *Thinking in Sound: The Cognitive Psychology of Human Audition* (pp. 157-213). Oxford: Oxford University Press.
- McAdams, S., Winsberg, S., Donnadieu, S., De Soete, G., & Krimphoff, J. (1995). Perceptual scaling of synthesized musical timbres: Common dimensions, specificities, and latent subject classes. *Psychological Research*, 58(3), 177-192.
- Moore, B. C. & Glasberg, B. R. (1983). Suggested formulae for calculating auditory-filter bandwidths and excitation patterns. *Journal of the Acoustical Society of America*, 74(3), 750-753.
- Opolko, F. & Wapnick, J. (1989). *McGill University Master Samples*. Montreal, QC: McGill University.
- Peeters, G., Giordano, B.L., Susini, P., Misdariis, N. & McAdams, S. (2011). The Timbre Toolbox: Extracting audio descriptors from musical signals. *Journal of the Acoustical Society of America*, 130, 2902-2916.
- Piston, W. (1955). *Orchestration*. New York: Norton.
- Reby, D., & McComb, K. (2003). Anatomical constraints generate honesty: acoustic cues to age and weight in roars of red deer stags. *Animal Behavior*, 65, 519-530.
- Rimsky-Korsakoff, N., & Steinberg, M. (1914). *Principes d'orchestration* (Vol. 1). Berlin: Édition russe de musique.
- Schaeffer, P. (1966). *Traité des objets musicaux : Essai interdisciplines*. Paris: Éditions du Seuil.
- Smith, D. R. R., Patterson, R. D., Turner, R., Kawahara, H., & Irino, T. (2005). The processing and perception of size information in speech sounds. *Journal of the Acoustical Society of America*, 117(1), 305-318.

- Štěpánek, J. (2006). Musical sound timbre: Verbal description and dimensions. In *Proc. of the Int. Conf. on Digital Audio Effects (DAFx-06), Montreal, Quebec, Canada* (pp. 121–126).
- Stevens, S. S. (1971). Issues in psychophysical measurement. *Psychological Review*, 75(5), 426–450.
- Stevens, S. S. (1975). *Psychophysics: Introduction to its Perceptual, Neural and Social Prospects*. New York: Wiley.
- Stevens, S. S., & Davis, H. (1938). *Hearing: Its Psychology and Physiology*. New York: Wiley.
- Stevens, S. S., Guirao, M., & Slawson, A. W. (1965). Loudness, a product of volume times density. *Journal of Experimental Psychology*, 69, 503–510.
- Stumpf, C. (1890). *Tonpsychologie [Tone Psychology]*, vol. 2. Leipzig: Hirzel.
- Terrace, H. S., & Stevens, S. S. (1962). The quantification of tonal volume. *American Journal of Psychology*, 75, 596–604.
- Traube, C., & Depalle, P. (2004). *Timbral analogies between vowels and plucked string tones*. Paper presented at the International Conference on Acoustics, Speech, and Signal Processing (ICASSP '04), Montreal, Canada.

TABLE Captions

- Table 1.** Volume scale of instruments of the orchestra given by Koechlin (1954-1959, vol. I, p. 288) and an example of the predicted scale according to Koechlin for the instruments used in the experiments on the pitch B4. We added solo strings, which were predicted by Koechlin to be the thinnest instruments.
- Table 2.** Series of instrument samples in Experiment 1 (columns) and their rankings according to Koechlin's volume scale. The first six series use a single pitch (F3-B5). The last two series use different pitches. Series 7 (TransA – light grey) uses instruments with predicted increasing extensity with increasing pitch, whereas Series 8 (TransB – dark grey) uses instruments with decreasing extensity with increasing pitch.
- Table 3.** Koechlin's volume scale and participants' ordinal extensity scales from both experiments. Instruments in the same squares are considered to have similar volumes, according to Bonferroni-Holm corrected significance values. As we see in Figures 2 and 3, if notches are not overlapping, we consider them to be in separate categories. Koechlin had himself separated and clustered instruments into categories. Instruments considered to have different ranks in Koechlin's and participants' scales are in italics.
- Table 4.** Rotated component matrix (Varimax rotation with Kaiser normalization) from the principal component analysis.

Table 1.


	Koechlin's scale	Experiment scale on B4
Large (<i>Gros</i>) 	Brass contrabasses, Tubas	
	Horns, Saxhorns	Horn
	Trombones	
	Trumpets, Flutes in low register, Bassoons, Saxophones	Trumpet Alto saxophone
	Clarinets, Oboe in low register, Flutes in medium register	Clarinet Flute
	Oboes in medium and high registers, Flutes in high register, Clarinets in altissimo register	Oboe
Thin (<i>Mince</i>)	<i>Solo strings</i>	Violin

Table 2.

Volume Rank	F3	B3	F4	B4	F5	B5
5		Horn	Horn	Horn	Horn	
4	Saxophone, Trumpet	Flute, Saxophone, Trumpet	Flute, Saxophone, Trumpet	Saxophone, Trumpet	Saxophone, Trumpet	Trumpet
3	Clarinet	Clarinet, Oboe	Clarinet, Oboe	Clarinet, Flute	Clarinet, Flute	Clarinet, Flute
2				Oboe	Oboe	Oboe
1		Violin	Violin	Violin	Violin	Violin
	Series 1	Series 2	Series 3	Series 4	Series 5	Series 6

Volume Rank	F3	B3	F4	B4	F5	B5
5					Horn	
4		Flute, Saxophone, Trumpet	Flute, Saxophone, Trumpet			
3		Clarinet, Oboe	Clarinet, (<i>Oboe not used</i>)			
2				Oboe		
1					Violin	
	Series 7 (TransA)				Series 8 (TransB)	

Table 3.

Koechlin's Volume Rank	F3			B3			F4		
	Koechlin	Exp. 1	Exp. 2	Koechlin	Exp. 1	Exp. 2	Koechlin	Exp. 1	Exp. 2
5				Horn	Sax Trumpet Horn	Sax Trumpet Horn	Horn	Sax Trumpet Horn	Horn Trumpet
4	Sax Trumpet	Sax	Sax	Flute Sax Trumpet			Flute Sax Trumpet		Sax
3	Clarinet	<i>Trumpet</i> Clarinet	<i>Trumpet</i> Clarinet	Clarinet Oboe	Clarinet	Clarinet Oboe	Clarinet Oboe	Clarinet	Clarinet
					Oboe			Oboe	<i>Flute</i> Oboe
2					Violin <i>Flute</i>	Violin <i>Flute</i>		<i>Flute</i>	
1				Violin			Violin	Violin	Violin

Koechlin's Volume Rank	B4			F5			B5		
	Koechlin	Exp. 1	Exp. 2	Koechlin	Exp. 1	Exp. 2	Koechlin	Exp. 1	Exp. 2
5	Horn	Sax Trumpet Horn	Trumpet Sax Horn	Horn	<i>Trumpet</i>	Trumpet Horn			
4	Sax Trumpet			Sax Trumpet	<i>Horn</i> Sax		Trumpet	Trumpet	Trumpet
3	Clarinet Flute	Clarinet Flute	Clarinet Flute	Clarinet Flute	Clarinet	Sax Clarinet Flute	Clarinet Flute	Clarinet	Clarinet
					Flute			Flute	Flute
2	Oboe	Violin Oboe	Oboe Violin	Oboe	Oboe	Oboe	Oboe	Oboe	Oboe
1	Violin			Violin	Violin	Violin	Violin	Violin	Violin

Table 3 (con't).

Koechlin's Volume Rank	TransA			TransB		
	Koechlin	Exp. 1	Exp. 2	Koechlin	Exp. 1	Exp. 2
5	Horn F5	Trumpet F4 Sax F4 Horn F5	Sax F4 Trumpet F4 Horn F5			
4	Flute F4 Sax F4 Trumpet F4			Flute B3 Sax B3 Trumpet B3	Trumpet B3 Sax B3	Sax B3 Trumpet B3
3	Clarinet B3 Oboe B3	Clarinet B3 Oboe B3	Clarinet B3 Oboe B3	Clarinet F4	Clarinet F4	Clarinet F4
2		<i>Flute F4</i>	<i>Flute F4</i>	Oboe B4	<i>Flute B3</i> Oboe B4 Violin F5	Oboe B4
1				Violin F5		<i>Flute B3</i> Violin F5

Table 4.

Acoustic descriptor	PC1	PC2	PC3
Kurtosis	−0.933	−0.044	−0.081
Spectral spread	0.915	0.224	0.063
Skewness	−0.914	−0.207	−0.119
Absolute spectral centroid	0.713	0.65	0.128
Loudness	−0.617	0.057	0.443
Spectral roll-off	0.14	0.892	0.107
Spectral irregularity	−0.085	−0.825	0.002
Fundamental frequency	0.236	0.24	0.85
Relative spectral centroid	0.297	0.188	−0.808
Attack time	0.435	0.498	−0.573
Spectral flux	0.015	0.352	0.491
Even/odd ratio	−0.156	−0.006	−0.451

FIGURE Captions

Figure 1. Diagram of relations among orchestration descriptors used in Koechlin's *Traité de l'orchestration* (translated from Chiasson, 2010).

Figure 2. Box-and-whiskers plots of participants' rankings of instrument largeness compared to Koechlin's volume scale for each set of Experiment 1. The x-axis gives Koechlin's ranking. The y-axis gives participants' rankings. The center line of each box indicates the median. The box indicates the interquartile range (IQR). Notches indicate $1.5 \cdot \text{IQR} / \sqrt{n}$ as a measure of confidence interval. Whiskers extend to $\pm 1.5 \cdot \text{IQR}$. Crosses indicate outliers. In some cases, the confidence interval goes beyond the quartiles.

Figure 3. Box-and-whiskers plots of participants' ratio estimates of instrument largeness (log scale) compared to Koechlin's ordinal volume scale for all sets. (See Figure 2 caption.)

Figure 4. Models of a large sound and a small sound according to the most salient acoustic features revealed by the Principal Components Analysis.

Figure 5. Harmonic spectra and spectral envelopes of instruments playing F4, in the order of Koechlin's volume scale : from left to right (violin to clarinet), then on the second row from left to right again (saxophone to horn).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

