

Nitrogen deposition and increased carbon accumulation in ombrotrophic peatlands in eastern Canada

Jukka Turunen,¹ Nigel T. Roulet,^{1,2} and Tim R. Moore¹

Department of Geography, McGill University, Montreal, Quebec, Canada

Pierre J. H. Richard¹

Département de Géographie, Université de Montréal, Montreal, Quebec, Canada

Received 23 September 2003; revised 25 March 2004; accepted 19 April 2004; published 10 July 2004.

[1] Recent and long-term accumulation rates of carbon (C), using ²¹⁰Pb- and ¹⁴C-dating, were examined in 23 ombrotrophic peatlands in eastern Canada, where average 1990–1996 atmospheric wet nitrogen (N) deposition ranged from 0.3 to 0.8 g N m⁻² yr⁻¹. The average recent rate of C accumulation (RERCA) over the past 150 years was 73 ± 17 (SD) g C m⁻² yr⁻¹, ranging from 40 to 117 g C m⁻² yr⁻¹. The difference in RERCA between hummocks (78 g C m⁻² yr⁻¹) and hollows (65 g C m⁻² yr⁻¹) was significant. Increased RERCA over the past 50 years was found in hummocks and hollows in regions of higher N deposition and related to both elevated N deposition and growing degree-days above +5°C. There was a statistically significant positive relationship between N deposition alone and present-day C accumulation in both hummocks and hollows (R² = 0.28 and 0.38, respectively). Recent N accumulation was significantly larger in high N deposition regions. The total average aboveground vegetation biomass of hollows and hummocks did not differ significantly with N deposition. However, a significantly larger vascular plant leaf biomass was found in both hollows and hummocks of the high N deposition class than in the low N deposition class (>0.6 and <0.4 g m⁻² yr⁻¹, respectively). The average long-term apparent rate of C accumulation (LORCA) at 15 sites was 19 ± 8 (SD) g C m⁻² yr⁻¹, with no significant difference due to age of peat inception, latitude, or continentality. *INDEX TERMS*: 4806 Oceanography: Biological and Chemical: Carbon cycling; 0315 Atmospheric Composition and Structure: Biosphere/atmosphere interactions; 1615 Global Change: Biogeochemical processes (4805); 1040 Geochemistry: Isotopic composition/chemistry; *KEYWORDS*: carbon accumulation, nitrogen, ombrotrophic peatlands, raised bogs, ²¹⁰Pb dating, Canada

Citation: Turunen, J., N. T. Roulet, T. R. Moore, and P. J. H. Richard (2004), Nitrogen deposition and increased carbon accumulation in ombrotrophic peatlands in eastern Canada, *Global Biogeochem. Cycles*, 18, GB3002, doi:10.1029/2003GB002154.

1. Introduction

[2] Global concern over rising atmospheric CO₂ concentration has led to attempts to ascertain the role of terrestrial ecosystems in the global carbon (C) cycle. Forests and peatlands in the Northern Hemisphere have been identified as potentially large sinks for organic C [Tans *et al.*, 1990; Gorham, 1991; Kauppi *et al.*, 1992; Houghton, 1993]. On a global scale, the occurrence of peatlands is strongly related to topography and climate, with the greatest abundance found on flat land areas of cool and moist climates, such as western Siberia, Russia, and the Hudson Bay Lowlands of Canada [Sjörs, 1959; Botch *et al.*, 1995].

Organic peat deposits are characterized by C contents of approximately 50% of dry matter [Gorham, 1991; Turunen *et al.*, 2001, 2002]. Therefore the abundance of peat signals a significant net transfer of C from the atmosphere to soils.

[3] Gorham [1991] has estimated the total area of boreal and subarctic peatlands to be 346 million ha and that 200 to 455 Pg C has accumulated during the Holocene [Gorham, 1990, 1991] with an update of 270 to 370 Pg C [Turunen *et al.*, 2002]. Despite the uncertainties in the storage estimates, peatlands are a substantial reservoir of C in the boreal and subarctic regions, constituting at least one fifth of the total soil C pool in the world [Post *et al.*, 1982], and being approximately half the amount of CO₂-C in the atmosphere [Houghton *et al.*, 1990].

[4] Peatlands generally accumulate C because the rate of biomass production is greater than the rate of decomposition. The average long-term apparent rate of carbon accumulation (LORCA) for boreal and subarctic regions throughout the Holocene is estimated at 15 to 30 g C m⁻² yr⁻¹

¹Also at Centre for Climate and Global Change Research, McGill University, Montreal, Quebec, Canada.

²Also at McGill School of Environment, McGill University, Montreal, Quebec, Canada.




Figure 1. Location and N deposition class of the 23 ombrotrophic peatlands sampled in eastern Canada. The aboveground vegetation biomass sampling sites are also shown.

[Gorham, 1990, 1991; Vitt *et al.*, 2000; Turunen *et al.*, 2001, 2002]. Eddy covariance and hydrological measurements at the Mer Bleue peatland, near Ottawa, Canada, indicate an annual sink of approximately 60 g C m^{-2} for the last 3 years [Lafleur *et al.*, 2001]: a rate 3 times greater than the long-term accumulation inferred from peat cores in the same peatland (P. J. H. Richard, unpublished data, 2004).

[5] There are a number of reasons why current Mer Bleue C accumulation rates may be higher: Climate variability, nitrogen deposition, and response to elevated CO_2 . CO_2 response should be universal and not isolated to eastern Canada. Recent studies have indicated that while plant C allocation changes in response to elevated CO_2 , there is no significant increase in overall peatland net primary production [Berendse *et al.*, 2001]. Long-term climate variability likely alters C accumulation on a decadal scale but is also spatially extensive, for example, climate variability associated with the North Atlantic/Arctic Oscillation affects most of northeastern North America. However, nitrogen (N) deposition varies across eastern North America and may affect C accumulation through effects on plant productivity and rates of organic matter decomposition. Mer Bleue is located in a region of some of the highest N deposition in eastern North America ($\sim 0.8 \text{ g N m}^{-2} \text{ yr}^{-1}$), while the Maritimes receive moderate levels of N deposition and the lowest rates are observed in northwestern Ontario ($\sim 0.3 \text{ g N m}^{-2} \text{ yr}^{-1}$).

[6] The specific objective of the research was to investigate the recent (RERCA) and long-term (LORCA) rates of carbon accumulation of 23 ombrotrophic bogs in eastern Canada and to test for relationships of these rates with N deposition and other environmental variables. Also, the recent rates of nitrogen accumulation (RERNA) were investigated.

2. Materials and Methods

2.1. Study Area and Sampling

[7] Twenty-three ombrotrophic bogs were selected (Figure 1, Table 1) based on the work of Gorham *et al.* [1985], Damman and Dowhan [1981], Glaser and Janssens [1986], and Damman [1988]. All peatlands were characterized by an open canopy of *Picea mariana* (approximately treeless), with *Chamaedaphne calyculata*, *Ledum groenlandicum*, *Kalmia angustifolia*, *Kalmia polifolia*, *Vaccinium uliginosum*, *Vaccinium myrtilloides*, *Vaccinium oxycoccos*, *Andromeda polifolia*, and scattered *Eriophorum spissum*. In coastal regions, the shrub cover was less dominant compared to more continental areas. However, the same species were found in all sampled bogs. A few *Pinus banksiana* and *Larix laricina* were found as a mixture with *Picea mariana* in several peatlands of Ontario and Québec. Hummocks were dominated by *Sphagnum fuscum*, with scattered *S. magellanicum*, *S. capillifolium*, *Polytrichum* spp., *Cladina* spp., *Empetrum nigrum*, and *Sarracenia purpurea*, and hollows by *S. rubellum* and *S. angustifolium*. In coastal regions, *S. flavicomans* was commonly found in the sampled hummocks as a mixture with *S. fuscum*. Scattered *Juniperus communis* was also found in the hummocks of peatlands near the Atlantic Ocean.

[8] The RERCA and RERNA were measured at all 23 bogs by collecting three short cores (50 cm) using a box sampler ($85 \times 85 \times 1000 \text{ mm}$), two from hummock sites and one from a hollow site. Each short core was double-wrapped in polythene bags, and stored at -4°C . Additional peat samples were collected down to 80 cm for each core in case they were needed for further analysis. The LORCA was measured at 15 bogs (Table 2). Complete peat profiles with contiguous 10-cm intervals were collected from hummock sites for dry bulk density measurements with the box sampler and a Russian pattern side-cutting peat sampler ($50 \times 500 \text{ mm}$). The degree of decomposition was estimated in the field with von Post's [1922] 10-grade scale (H_{1-10}). Basal peat samples from the coring sites were taken for ^{14}C dating. The aboveground vegetation was collected from 0.25 m^2 plots at two or more hummock sites and one hollow site at 13 bogs (Figure 1).

2.2. Laboratory Analyses

[9] The peat samples were dried to a constant mass at 70°C and weighed and the dry bulk density calculated by dividing the dry mass by the fresh volume. The C and N concentrations of peat were analyzed using an Elemental vario EL analyzer at University of Bayreuth, Germany [Tabatabai and Bremer, 1990]. The cumulative dry mass of peat on an areal basis (g m^{-2}) was calculated as layer thickness weighted averages from the dry bulk density profile and converted into C and N, based on C and N concentration analyses for surface cores (max top 80 cm) and 51.7% of C for long cores [Gorham, 1991]. The accumulated mass of C and N above the oldest ^{210}Pb -datable horizon was divided by the age of this horizon to give the RERCA ($\text{g C m}^{-2} \text{ yr}^{-1}$) and RERNA ($\text{g N m}^{-2} \text{ yr}^{-1}$). The LORCA was determined similarly, but using the C mass above the peat-mineral contact, and the age of the basal peat sample. The Mer Bleue Bog was the exception, in that the

Table 1. Location and Characteristics of Ombrotrophic Peatlands Sampled in Canada (See Also Figure 1) and Characteristics of the Sites in the Three N Deposition Classes^a

N Class	Peatland	North Latitude	West Longitude	Mean Temperature, °C			Effective Temperature Sum, dd	Annual Precipitation			N Deposition, g m ⁻² yr ⁻¹	
				January	July	Annual		Rain, mm	Snow, cm	Total, mm		
1	1 Petite Bog, NS	45°09'	63°56'	-5.7	17.1	6.5	1602	944	230	1175	0.34 (0.08)	
	2 Western Head, NS	43°41'	65°08'	-1.6	11.3	5.9	1000	1068	149	1217	0.33 (0.04)	
	3 Cape Sable, NS	43°28'	65°36'	-1.6	11.3	5.9	1000	1068	149	1217	0.33 (0.04)	
	4 Fourchu, NS	45°42'	60°14'	-5.4	17.6	5.5	1444	1156	330	1480	0.29 (0.03)	
	5 Foxley Moor, P.E.I.	46°43'	64°02'	-8.6	18.6	5.0	1545	812	278	1090	0.32 (0.08)	
	6 Point Sapin, NB	46°59'	64°51'	-10.4	19.1	4.6	1591	768	324	1087	0.32 (0.08)	
	7 Point Escuminac, NB	47°04'	64°49'	-10.4	19.1	4.6	1591	768	324	1087	0.27 (0.06)	
	8 Miscou Island, NB	47°56'	64°30'	-10.4	19.1	4.6	1611	710	308	1018	0.28 (0.07)	
	9 Savoy Bog, NB	47°47'	64°36'	-10.4	19.1	4.6	1611	710	308	1018	0.28 (0.07)	
	11 Yellow Lake, Qc	48°54'	71°54'	-17.0	17.3	1.5	1363	637	197	835	0.32 (0.06)	
	14 Ilets-Jeremie Bog, Qc	48°54'	68°49'	-14.0	15.6	1.5	1026	662	362	996	0.35 (0.06)	
	15 Mai Bog, Qc	49°57'	67°02'	-14.6	15.2	1.0	947	729	415	1128	0.33 (0.05)	
	16 Port Cartier Bog, Qc	50°02'	66°56'	-14.6	15.2	1.0	947	729	415	1128	0.29 (0.05)	
	17 Escumins Bog, Qc	48°24'	69°21'	-12.5	16.9	3.0	1284	691	310	998	0.36 (0.05)	
	19 Baker Bog, Ont	49°08'	90°45'	-17.9	17.7	1.5	1338	581	238	763	0.35 (0.06)	
	Mean				-10.2 (4.9)	16.6 (2.5)	3.8 (1.9)	1320 (269)	808 (171)	289 (83)	1090 (160)	0.32 (0.03)
	2	13 Despinassy Bog, Qc	48°44'	77°44'	-14.3	16.1	1.0	1141	670	244	913	0.47 (0.10)
		20 Norembego, Ont	48°59'	80°42'	-18.2	16.7	0.6	1206	604	316	920	0.52 (0.12)
		21 Nellie Bog, Ont	48°46'	80°50'	-17.6	17.1	0.9	1252	573	219	793	0.52 (0.12)
22 Hislop Bog, Ont		48°28'	80°23'	-17.1	17.7	1.5	1335	587	289	876	0.52 (0.12)	
23 Holtrye Bog, Ont		48°32'	80°48'	-17.6	17.1	0.9	1252	573	219	793	0.52 (0.12)	
Mean					-17.4 (1.0)	17.0 (0.4)	0.9 (0.3)	1245 (54)	591 (28)	257 (43)	848 (60)	0.51 (0.02)
3	10 Lac à la Tortue, Qc	46°32'	72°40'	-12.7	19.6	4.6	1738	783	250	1042	0.76 (0.14)	
	12 Mirabel, Qc	45°41'	74°03'	-11.7	19.7	4.8	1748	796	233	1030	0.81 (0.12)	
	18 Mer Bleue, Ont	45°25'	75°31'	-10.8	20.8	5.8	1941	702	222	910	0.81 (0.13)	
	Mean				-11.7 (1.0)	20.0 (0.7)	5.1 (0.6)	1821 (109)	753 (47)	235 (14)	987 (70)	0.79 (0.02)

^aClass 1: <0.4; class 2: 0.4–0.6; class 3: >0.6 g N m⁻² yr⁻¹. For effective temperature sum, a +5°C threshold is used, dd is degree-days. Standard deviation (SD) in parentheses. NS, Nova Scotia; P.E.I., Prince Edward Island; NB, New Brunswick; Qc, Québec; Ont, Ontario.

basal 70 cm of the limnic peat was excluded from the LORCA analysis, and the pond-fen transition zone (500 cm) was used as a basal peat depth.

[10] Ten samples of 1 cm thickness were taken from each peat core at 5-cm intervals and submitted to Flett Research Ltd., Winnipeg, Canada for ²¹⁰Pb dating. Similarly, extra samples from 50 to 80 cm, especially from hummock sites, were also analyzed if the unsupported ²¹⁰Pb activity was much greater than the supported ²¹⁰Pb activity (i.e., measured activity had not reached the low natural ²¹⁰Pb background activity). The theory of ²¹⁰Pb-dating ($t_{1/2} =$

22.3 years) assumes that ²¹⁰Pb concentrations in peat decrease exponentially with depth, approaching a low constant value taken to represent the supported ²¹⁰Pb fraction formed within soil as opposed to that deposited from the atmosphere [e.g., *Turetsky et al.*, 2000]. A dry 0.5-g subsample from each peat layer was analyzed for ²¹⁰Pb activity (as estimated by the alpha emitting ²¹⁰Po granddaughter), after spiking with a ²⁰⁹Po yield tracer. The constant rate of supply (CRS) model of *Appleby and Oldfield* [1978] was applied to calculate the ages of peat layers. The C accumulation of the last 50 years was used as

Table 2. Basal Sample Characteristics and the Results of ¹⁴C Dating in Ombrotrophic Peatlands of Eastern Canada^a

Site	Sample Depth, cm	Peat Material	Lab. Number	¹³ C/ ¹² C	¹⁴ C Date, Year BP	Calendar Date, Year BP	LORCA, g C m ⁻² yr ⁻¹
1 Petite Bog, NS	340–345	L-C, H5	Beta-151184	-26.8	3450 ± 70	3750	27.8
2 Western Head, NS	585–590	C, H6-7	Beta-151185	-27.8	7970 ± 120	8920	18.5
4 Fourchu, NS	145–150	N-Er-C, H4-5	Beta-151186	-27.1	2380 ± 50	2410	16.1
5 Foxley Moor, P.E.I.	105–110	Pr-C, H7	Beta-151187	-25.5	7840 ± 70	8660	5.8
6 Point Sapin, NB	340–345	Eq-C, H9	Beta-151188	-25.7	6460 ± 70	7460	13.2
7 Point Escuminac, NB	495–500	N-S-C, H9	Beta-151189	-27.5	4170 ± 60	4800	28.6
8 Miscou Island, NB	630–635	N-S, H4-5	Beta-151190	-27.1	8070 ± 100	9060	22.9
10 Lac à la Tortue, Qc	415–420	C, H9	Beta-151191	-28.0	8150 ± 70	9090	16.6
11 Yellow Lake, Qc	290–295	C, H7-8	Beta-151192	-28.8	5420 ± 60	6290	18.0
13 Despinassy Bog, Qc	145–150	C-B, H4	Beta-151193	-26.9	6760 ± 70	7650	5.1
14 Ilets-Jeremie Bog, Qc	370–375	N-Sch-S-C, H5	Beta-151194	-26.9	3570 ± 60	3910	34.6
16 Port Cartier Bog, Qc	265–270	N-S, H8	Beta-151195	-26.8	5750 ± 70	6580	17.3
18 Mer Bleue, Ont	475–480	N-S-C, H5	TO-8163	-	7340 ± 70	8170	24.3
19 Baker Bog, Ont	215–220	Sch-C, H5	Beta-151196	-26.3	8700 ± 80	9710	10.0
20 Norembego, Ont	320–325	L-S, H6-7	Beta-151197	-26.7	5540 ± 50	6360	23.5

^aLORCA, the long-term apparent rate of carbon accumulation. Peat constituents: B, Bryales; C, *Carex*; Er, *Eriophorum*; Eq, *Equisetum*; L, wood; N, shrub; Pr, *Phragmites*; S, *Sphagnum*; Sch, *Scheuchzeria*; H, degree of decomposition in *von Post's* [1922] ten-grade scale. Calendar date refers to the calibrated calendar years [*Stuiver and Reimer*, 1993].

the basis mass for C loss comparison because ^{210}Pb -dating has proven most reliable for peat deposits in the range of 50–200 years [Malmer and Holm, 1984].

[11] Basal peat samples were ^{14}C dated at the Beta Analytical Dating Laboratory, Florida, and University of Toronto, Ontario, Canada. All results were corrected for isotopic fractionation based on the ^{13}C -values (Table 2) and radiocarbon ages were converted to calendar years using CALIB 3.0.3 [Stuiver and Reimer, 1993]. All ages used in this article are calendar years (cal. BP).

[12] Aboveground biomass was separated into vascular leaves and stems and *Sphagnum capitulum*, dried to a constant mass at 70°C and weighed [Moore et al., 2002].

[13] The climate data (Table 1) were derived from the Canadian Climate Normals (Meteorological Service of Canada, http://www.msc-smc.ec.gc.ca/climate/climate_normals/index_e.cfm), using the station nearest each bog. The wet N deposition ($\text{N-NH}_4 + \text{N-NO}_3$) was calculated using the 7-year mean (1990–1996) for eastern North America [Vet et al., 1999], using the station nearest each bog. The bogs were allocated to three N deposition classes: class 1: <0.4 , class 2: $0.4\text{--}0.6$, class 3: $>0.6\text{ g N m}^{-2}\text{ yr}^{-1}$ (Table 1).

[14] SPSS for Windows 10.0 statistical software was used to analyze the relationships between variables. The normal distribution and homogeneity of variances were tested using the Kolmogorov-Smirnov and Levene statistical tests. Stepwise, ordinary least squares multiple regression was used to examine which characteristics of the sites (Table 1) were related to the rates of C accumulation. Pearson's correlation coefficient r was used for correlation analysis. To compare differences in the C accumulation and in biomass of hollows and hummocks between study sites along the atmospheric N deposition gradient, analysis of variance (ANOVA) was conducted.

3. Results

3.1. ^{210}Pb Concentrations

[15] Mean total residual unsupported (i.e., Pb deposited by atmospheric processes as opposed to Pb incorporated into the biomass) ^{210}Pb activity in the surface cores was $0.35 \pm 0.11\text{ Bq cm}^{-2}$ (counting time 500 min), with larger mean values in hummocks (0.38 Bq cm^{-2}) than hollows (0.30 Bq cm^{-2}). The geographic pattern of unsupported ^{210}Pb burden in peat was similar to that of industrial N deposition. In N deposition classes 1, 2, and 3, the average unsupported ^{210}Pb burden in hollows was 0.25, 0.39, and 0.42 Bq cm^{-2} and in hummocks 0.37, 0.38, and 0.45 Bq cm^{-2} , respectively. The ^{210}Pb -dating worked well in all 23 hollows but difficulties arose in some hummocks in the high N-deposition class. These hummocks were characterized by a large shrub biomass and weakly decomposed, loose *Sphagnum fuscum*. Eight hummocks were subsequently excluded because data inconsistencies in ^{210}Pb -activity throughout the peat columns made it unreliable to construct ^{210}Pb -based chronologies for these cores.

3.2. Short-Term C Accumulation

[16] The RERCA over the past 150 years revealed considerable variation among different peatlands, ranging from

40 to $117\text{ g C m}^{-2}\text{ yr}^{-1}$ (Table 3). Results from site # 19 were excluded from the analysis as outliers (Table 3). The average dry bulk density and C concentration among hummocks (42 g dm^{-3} , 46%) and hollows (51 g dm^{-3} , 46%) were similar. RERCA was significantly larger in hummocks than in hollows over the past 150 years. Also, the variation in RERCA and cumulative C mass within hummocks was much less than within hollows (Table 3, Figure 2). Generally, the average net accumulation rate of C decreased with time both in hollows and hummocks because of decay, even though some variation was found among peatlands.

[17] The effect of climate characteristics (Table 1) on RERCA was investigated by stepwise, ordinary least squares multiple regressions (entry level 0.10). Since the increase in N deposition has been mainly in the second half of the twentieth century, the 50-year RERCA was used in this analysis. Multiple regressions revealed significant relationships between recent C accumulation, climate characteristics, and the N deposition. The 50-year RERCA rates in hollows and hummocks were best predicted from N deposition and growing degree-days above $+5^\circ\text{C}$ ($r^2 = 0.46$ and 0.36 , respectively; Figure 3, Table 4). Generally, multicollinearity between climate characteristics was a problem. However, the predictive variables in both regressions were relatively independent ($r = 0.44$ and $r = 0.31$, respectively). Overall, the relatively weak relationships between climate variables and C accumulation over the 100 and 150 years were very similar to each other. The 150-year RERCA rates in hollows and hummocks were best predicted from annual total precipitation (negative, $r^2 = 0.18$ and 0.26 , respectively).

[18] Analysis of variance (ANOVA) revealed that RERCA was significantly greater in hummocks than in hollows in N deposition classes 1 and 2 over the last 50, 100, and 150 years. Differences in RERCA within high N deposition class 3, however, were not statistically significant over the last 50, 100, and 150 years. Differences in RERCA between classes of different rates of N deposition were significant. The 50-year RERCA was significantly larger in hollows of the high N deposition class 3 compared to hollows in lower N deposition classes 1 and 2 ($p = 0.004$ and $p = 0.030$, respectively). The 100- and 150-year RERCA were significantly larger in hollows of high N deposition class 3 compared to low N deposition class 1 ($p = 0.005$ and $p = 0.007$, respectively). In hummocks, the 50-year RERCA was significantly larger in the high N deposition class 3 compared to lower N deposition class 1 ($p = 0.027$). The 50-year RERCA was significantly larger in the N deposition class 2 compared to lower N deposition class 1 ($p = 0.012$), though differences in 100- and 150-year RERCA were not statistically significant, but the trend was to an increasing RERCA toward the higher N deposition classes (Table 3).

[19] Assuming that C input into the acrotelm, the upper oxic zone of the peat profile, has been relatively constant during the last 150 years, the hummock and hollow peat layers from 50 to 100 years have lost approximately 38% and peat layers from 100 to 150 years have lost approximately 57% of the original C mass compared to the recent

Table 3. Recent Apparent Rate of Carbon Accumulation (RERCA) Based on ^{210}Pb Dating in Ombrotrophic Peatlands of Eastern Canada^a

N Class	Peatland	Hollows		RERCA, g C m ⁻² yr ⁻¹		Hummocks		
		50 Years	100 Years	~150 Years	50 Years	100 Years	~150 Years	
1	1 Petite Bog, NS	83.5	68.4	58.2	77.1 (15.3)	64.4 (3.6)	61.2 (5.7)	
	2 Western Head, NS	68.9	48.6	49.7	128.7 (35.2)	90.6 (-)	78.8 (-)	
	3 Cape Sable, NS	85.0	69.1	64.8	101.3 (1.5)	79.3 (5.2)	-	
	4 Fourchu, NS	67.2	52.3	43.8	109.7 (6.0)	77.6 (14.6)	61.8 (10.4)	
	5 Foxley Moor, P.E.I.	97.5	92.8	81.3	115.1 (9.9)	113.1 (51.6)	92.5 (35.1)	
	6 Point Sapin, NB	68.6	53.3	45.3	128.2 (-)	98.1 (-)	-	
	7 Point Escuminac, NB	108.3	83.3	70.8	107.6 (5.4)	84.2 (14.3)	72.8 (4.9)	
	8 Miscou Island, NB	112.6	66.9	49.3	104.0 (4.8)	86.4 (2.9)	73.9 (0.1)	
	9 Savoy Bog, NB	54.0	40.3	42.4	101.6 (1.2)	89.9 (6.8)	74.0 (9.3)	
	11 Yellow Lake, Qc	84.1	79.8	68.9	99.5 (5.2)	83.7 (2.9)	71.0 (5.2)	
	14 Ilets-Jeremie Bog, Qc	83.9	71.2	56.7	126.4 (4.3)	94.1 (1.8)	79.3 (1.6)	
	15 Mai Bog, Qc	32.5	69.9	68.4	102.0 (15.7)	83.2 (9.2)	68.4 (6.4)	
	16 Port Cartier Bog, Qc	50.5	50.9	39.9	121.2 (3.4)	94.2 (11.9)	76.3 (9.0)	
	17 Escoumins Bog, Qc	130.4	92.5	76.9	128.6 (12.8)	104.7 (6.1)	84.9 (1.1)	
	19 Baker Bog, Ont	99.9	128.0	-	178.1 (-)	140.5 (-)	121.4 (-)	
	Mean	80.5 (26.3)	67.1 (16.4)	58.3 (13.6)	110.1 (17.0)	88.3 (16.9)	74.2 (12.1)	
	2	13 Despinassy Bog, Qc	111.9	105.7	86.0	-	-	-
		20 Norembego, Ont	92.6	83.8	76.5	135.1 (13.7)	102.8 (5.5)	86.7 (2.1)
		21 Nellie Bog, Ont	96.0	69.5	55.2	140.0 (20.2)	115.4 (0.8)	102.7 (4.2)
22 Hislop Bog, Ont		59.4	59.6	50.9	125.3 (-)	89.6 (-)	78.7 (-)	
23 Holtrye Bog, Ont		81.7	69.3	67.5	115.0 (11.5)	96.1 (7.2)	82.5 (6.8)	
Mean		88.3 (19.4)	77.6 (17.9)	67.2 (14.6)	131.7 (17.7)	102.6 (10.5)	88.9 (10.4)	
3	10 Lac à la Tortue, Qc	117.8	89.7	75.7	138.1 (-)	116.4 (-)	90.9 (-)	
	12 Mirabel, Qc	179.6	139.2	116.1	141.2 (-)	95.5 (-)	-	
	18 Mer Bleue, Ont	126.2	96.3	81.4	133.5 (-)	104.1 (-)	89.4 (-)	
	Mean	141.2 (33.5)	108.4 (26.9)	91.0 (21.9)	137.7 (3.9)	105.4 (10.7)	90.2 (1.1)	

^aStandard deviation (SD) in parentheses. Results from site 19 were excluded from the analysis as outliers. N deposition classes are as in Table 1.

50-year net C accumulation (Figure 4). The corresponding C loss values were greater for hummocks (40 and 59%) than hollows (34 and 52%) during the last 100 and 150 years, respectively. There was a uniform relative C mass loss of hollows among different N deposition classes; that is, the remaining total C mass increased consistently toward the highest N deposition class (Figure 4). A similar trend was seen in hummocks over the last 50 years; however, in older hummocks the differences between N deposition classes were marginal.

3.3. Long-Term C Accumulation

[20] The long-term apparent rate of carbon accumulation (LORCA) at 15 sites ranged from 5 to 34 g C m⁻² yr⁻¹ with an average of 19 ± 8 (SD) g C m⁻² yr⁻¹ (Figure 5, Table 2). LORCA values for sites 5 and 13 were exceptionally small (average 5.5 g C m⁻² yr⁻¹) compared to those of the other study sites (average 20.9 g C m⁻² yr⁻¹). In site 5 (depth 110 cm), there were several charcoal layers at short intervals between 72 and 88 cm, suggesting that the shallow peat layers had burned often, and in site 13 (depth 150 cm), thick charcoal horizons were found at 50 to 67 cm. No such evidence of peatland fires was found at the other sites. The effect of climate characteristics (Table 1) on LORCA was investigated by stepwise, ordinary least squares multiple regressions. Regression revealed no significant relationships between long-term C accumulation and present-day climate characteristics. Also, differences in LORCA between present-day N deposition classes 1, 2, and 3 were not significant ($p = 0.96$). There was a weak relationship between LORCA and RERCA (excluding sites

5, 13 and 19, Pearson's correlation coefficient $r = -0.01 - 0.22$, $p = 0.49 - 0.98$) over the last 50–150 years. Overall, the average RERCA:LORCA ratio for peat deposits of 50, 100, and 150 years was 8:1, 6:1, and 5:1, respectively.

3.4. Short-Term N Accumulation

[21] The mean N concentration of the samples was 0.82%, ranging from 0.35 to 2.25% with the hollows having a slightly larger average N concentration (0.90%) than the hummocks (0.76%). N concentrations were significantly larger in hollows of the higher N deposition classes 2 and 3 than hollows of low N deposition class 1 ($p = 0.001$ and $p = 0.001$, respectively). In hummocks, N concentrations were significantly larger in the higher N deposition classes 2 and 3 than hummocks of low N deposition class 1 ($p = 0.006$ and $p = 0.032$, respectively).

[22] The recent rate of nitrogen accumulation (RERNA) decreased with time in both hollows and hummocks. Average RERNA rates over the past 50, 100, and 150 years were 1.8, 1.7, and 1.5 g N m⁻² yr⁻¹, respectively. There were significant differences in RERNA among hollows and hummocks in classes of different rates of N deposition (Figure 6). The 50-year RERNA was significantly larger in hollows of the high N deposition class 3 than the lower N deposition class 1 ($p = 0.004$), but differences between N deposition classes 2 and 3 were not statistically significant ($p = 0.090$). However, the remaining N mass of peat layers from 100 to 150 years in the high N deposition class 3 was not significantly larger than that in the lower N deposition class 1 ($p = 0.116$). In hummocks, 50-year RERNA was significantly larger in higher N deposition classes 2 and 3




Figure 2. Cumulative C mass beneath the peat surface (kg C m^{-2}) versus year for (a) hollows ($n = 22$) and (b) hummocks ($n = 36$). Results from site 19 were excluded from the analysis as extreme outliers.

than low N deposition class 1 ($p = 0.001$ and $p = 0.027$, respectively). As in hollows, the remaining N mass of hummocks from 100 to 150 years in the high N deposition class 3 was not significantly larger than that in the lower N deposition class 1 ($p = 0.593$).

3.5. Aboveground Biomass

[23] Aboveground biomass was significantly greater on hummocks (633 g m^{-2}) than in hollows (482 g m^{-2}) (Table 5). Differences within different N deposition classes were evident, with the hummocks and hollows having a similar average *Sphagnum* capitula biomass within each N deposition class but a significantly greater vascular plant biomass being present on hummocks than in hollows (Table 5). The ANOVA did not reveal general, significant differences in the total average aboveground vegetation biomass of hollows and hummocks among N deposition classes. The trend, however, was toward smaller total above-

ground biomass in higher N deposition classes (Table 5). The only statistically significant difference found was with the smaller total average aboveground biomass on the hummocks of the highest N deposition class 3 compared to the lowest N deposition class 1. The general decrease of the total average aboveground biomass seemed to occur in the plant stem and *Sphagnum* capitula biomass. However, a significantly larger vascular plant leaf biomass was found both in hollows and hummocks of the highest N deposition class 3 than in classes 1 and 2 (Table 5). We also used stepwise, ordinary least squares multiple regression to examine if the total average aboveground biomass was related to local climate variables (Table 1) and/or N deposition over the last 50 years. The strongest predictor of aboveground biomass in the hummocks was N deposition ($r^2 = 0.41$) and in the hollows was total precipitation ($r^2 = 0.24$).

4. Discussion and Conclusions

4.1. ^{210}Pb Dating

[24] Mean total residual (unsupported) ^{210}Pb activity in the surface peat cores was 0.35 Bq cm^{-2} , well within the range of other soil inventories reported in North America ($0.31\text{--}0.84 \text{ Bq cm}^{-2}$ [Urban *et al.*, 1990]). We excluded eight hummocks because of inconsistencies in ^{210}Pb -activity throughout the peat cores, which may have resulted from fires or differential Pb deposition through the stems of shrubs, allowing partial wet and dry Pb deposition into deeper peat layers instead of *Sphagnum* capitula. No other disturbances, such as peat removal, had occurred in the sampling sites. The differences in the ^{210}Pb -activity of the surface moss layers and total ^{210}Pb burden in hollows and hummocks suggest that there must be differential deposition and/or post-depositional Pb mobility in some microhabitat types [Damman, 1978; Oldfield *et al.*, 1979; Pakarinen *et al.*, 1983; Urban *et al.*, 1990; Oldfield *et al.*, 1995]. However, several studies have concluded that Pb remains immobile after deposition on a peat surface [e.g., Livett *et al.*, 1979; Mitchell *et al.*, 1992; Appleby *et al.*, 1997; Vile *et al.*, 1999]. In this study, evidence of possible Pb mobility was also found in three hollows that showed exceptionally high ^{210}Pb -activity in the surface moss layers, suggesting post-depositional mobility from hummock to hollow or from depth to the surface. Mobility of ^{210}Pb seems to occur in some microhabitats, depending on different depositional, post-depositional, and geochemical conditions, such as distribution patterns of vegetation and regional differences in precipitation and peatland hydrology.

[25] As a check on the ^{210}Pb method, at Holtrye bog we were able to compare the ^{210}Pb -dates against a conspicuous charcoal horizon from a 1916 fire as a stratigraphical marker. In 1916, a fire in the Matheson area burned the villages of Kelso, Val Gagne, Porquis Junction, and Iroquois Falls, spreading over 500,000 ha and taking 244 lives (Ontario Ministry of Natural Resources, 2001, available at <http://www.mnr.gov.on.ca/MNR/affmb/Fire/Index/About/Default.htm>). The age difference in hollows and hummocks between the fire and ^{210}Pb -dating method was 4 and 8 years, respectively. Urban *et al.* [1990] suggested that hummocks are generally more reliable than hollows for preservation of historical records of atmospheric Pb deposition, and that age




Figure 3. Partial regression plots showing the influence of degree-days above °C and N deposition, on 50-year RERCA A–B) in hollows (n = 22), and C–D) in hummocks (n = 36), while removing the influence of the other independent variables. Results from site # 19 were excluded from the analysis as extreme outliers.

resolutions no better than ± 10 years may be attained with the ²¹⁰Pb-dating technique. Given the possible mobility of Pb, the interpretation of depth-age relationships in peat chronologies via ²¹⁰Pb-dating has to be made with caution, and data inconsistencies should be taken into account. However, a good agreement between ²¹⁰Pb-dates in peat and other dating methods suggests that the ²¹⁰Pb-dating is reliable over the past 200 years where the surface peat is too young for radiocarbon dating [El-Daoushy *et al.*, 1982; Belyea and Warner, 1994; Wieder *et al.*, 1994; Vile *et al.*, 1995; Appleby *et al.*, 1997; Shotyk *et al.*, 1997; Turetsky *et al.*, 2000].

4.2. Carbon Accumulation

[26] The C accumulation rate in eastern Canadian ombrotrophic bogs over the last 150 years ranged from 40

to 117 g C m⁻² yr⁻¹, with an average of 73 g C m⁻² yr⁻¹. This is similar to results from Finland (40 to 81 g C m⁻² yr⁻¹) [Tolonen and Turunen, 1996; Pitkänen *et al.*, 1999] and for boreal *Sphagnum* dominated peat deposits in North America (31–93 g C m⁻² yr⁻¹) [Tolonen *et al.*, 1988; Wieder *et al.*, 1994; Turetsky *et al.*, 2000]. Considerable variation in C mass accumulation was found both among and within peatlands, and differences in 150-year RERCA between hummocks and hollows were reflected in the average vertical height growth rates (of 4.0 and 2.8 mm yr⁻¹, respectively). The average vertical height growth rates in hummocks and hollows over the last 50 years were even more distinct (8.2 and 4.9 mm yr⁻¹, respectively). Økland and Ohlson [1998] reported a similar significant difference in vertical height growth between hummocks and hollows over the last 40 years in Scandinavian *Sphagnum* surface

Table 4. Regressions of the Recent Apparent Rate of Carbon Accumulation (RERCA) in Hollows and Hummocks Over the Last 50 Years^a

Equation	d.f.	F	P	r ²	s.e. _{est}
Fifty-year RERCA rates in hollows, g C m ⁻² yr ⁻¹					
RERCA = 4.20 + 8.91(N deposition) + 0.035 (dd)	1, 20	8.2	0.003	0.46	24.9
50-year RERCA rates in hummocks, g C m ⁻² yr ⁻¹					
RERCA = 111.21 + 8.18(N deposition) – 0.020(dd)	2, 34	9.4	0.001	0.36	16.0

^aNotation: dd, growing degree-days above +5°C; N deposition (g m⁻² yr⁻¹), annual mean temperature (°C), s.e._{est}, standard error of the estimate.




Figure 4. Total C mass in (a) hollows and (b) hummocks between 0 and 50 years, 50 and 100 years, and 100 and 150 years in three N deposition classes: class 1: <0.4 , class 2: $0.4\text{--}0.6$, class 3: >0.6 $\text{g N m}^{-2} \text{yr}^{-1}$. Vertical bars are standard errors.

peat. One reason for the RERCA/height growth differences is likely the faster decomposition rate in hollows than hummocks, resulting in larger dry bulk density values. The decomposition rates seem to be largest close to the water table [Belyea, 1996], and has been attributed to optimum conditions of aeration, water, and nutrient availability. Generally, the mean differences in *Sphagnum* capitulum biomass between hollows and hummocks were marginal (Table 5). However, comparisons are complicated because of a general substantial between-year variation in *Sphagnum* productivity within the same site [e.g., Lindholm and Vasander, 1990]. A significantly larger vascular plant biomass was found in hummocks than in hollows, which may cause variation in the autogenic factors of these microsites such as temperature and evapotranspiration and thus affect the recent C accumulation. There are indications that the fine-scale differences in the hydrology of each peatland




Figure 5. Average long-term apparent rate of carbon accumulation (LORCA) in ombrotrophic peatlands of eastern Canada compared to Finnish bogs [Tolonen and Turunen, 1996; Turunen et al., 2002]. The upper and lower limits of the range of LORCA in Finnish peatlands are also shown.

may play an important role in the observed variations of the aboveground biomass and the present-day C uptake [e.g., Ohlson and Dahlberg, 1991].

[27] The results of our study show that the age of the peat column is an important predictor of C accumulation rate (Figures 2 and 5, Table 3). The extensive data of Tolonen and Turunen [1996] also showed that the rate of C accumulation is considerably greater in younger peat deposits and the average net accumulation rate of C decreases with time because slow decay takes place in the anoxic, deeper peat layers [Clymo, 1984; Gorham, 1991; Tolonen and




Figure 6. Recent rate of nitrogen accumulation (RERNA) over the last 50 years in hollows and hummocks in ombrotrophic peatlands of eastern Canada.

Table 5. Average Aboveground Vegetation Biomass of Hollows and Hummocks in Ombrotrophic Peatlands in Eastern Canada^a

Microsite	n	Vascular Leaves, g m ⁻²	Vascular Stems, g m ⁻²	<i>Sphagnum</i> Capitulum, g m ⁻²	Total Biomass, g m ⁻²
<i>N Deposition Class 1</i>					
Hollows	5	95 (48) ^b	272 (241)	226 (94)	593 (331)
Hummocks	10	150 (30)	385 (184)	190 (42)	725 (190)
<i>N Deposition Class 2</i>					
Hollows	5	73 (24) ^b	196 (95) ^b	162 (36)	430 (117) ^b
Hummocks	10	144 (32)	279 (73)	180 (31)	603 (78)
<i>N Deposition Class 3</i>					
Hollows	4	162 (19) ^{c1-2}	114 (62) ^b	131 (62)	407 (111) ^b
Hummocks	8	180 (23) ^{c2}	233 (96) ^{c1}	142 (43) ^{c1}	555 (63) ^{c1}
<i>Mean</i>					
Mean hollows	14	106 (49) ^b	200 (161) ^b	176 (75)	482 (220) ^b
Mean hummocks	28	156 (32)	304 (140)	172 (43)	633 (143)
Total mean	42	139 (45)	269 (154)	174 (55)	582 (184)

^aStandard deviation (SD) in parenthesis. The N deposition classes as in Table 1.

^bSignificant difference between hollows and hummocks within N deposition class.

^cSignificant difference between hollows or hummocks among N deposition classes ($p < 0.05$). The index number after the footnote letter refers to the N deposition classes.

Turunen, 1996; Clymo *et al.*, 1998]. However, the 50-, 100-, and 150-year RERCA of both hollows and hummocks was not related to the age of the basal peat in this study (Tables 2 and 3); that is, the recent C accumulation was not less in older bogs than in younger ones. Our sample size, however, is relatively small, with only three bogs having a basal age younger than 4000 years, so the effect of the total peatland age or development stage on recent C accumulation remains uncertain.

[28] We have shown a statistically significant relationship between recent C accumulation, N deposition, and climate variables. The highest RERCA during the last 50 years occurred in the high atmospheric N deposition region, and the reason for this elevated RERCA seemed to be the combination of climate and high N deposition. Contrary to the results of Økland and Ohlson [1998] in Scandinavian surface peat, the among-peatland variations in C accumulation and in peat growth rates in eastern Canada were related to climate variables. Supporting our results, a rather clear relationship between long-term C accumulation and climate characteristics (degree-days and mean annual temperature) has been found for peatland regions in Finland and the mid-Boreal region of Canada [Clymo *et al.*, 1998].

[29] The increased C accumulation in ombrotrophic hummocks and hollows may be associated with N as a limiting element in most boreal ecosystems [Malmer, 1990; Aerts *et al.*, 1992]. Wet and dry N deposition, together with N fixation, are the only N sources of ombrotrophic peatland systems [Vaughman and Bellamy, 1980; Chapman and Hemond, 1982]. N fixation is probably less on hummocks than in the wetter hollows and may increase with increasing temperature and humidity [Urban and Eisenreich, 1988]. Increased C accumulation can also result from a decrease in decomposition rates. Some studies have indicated an increased accumulation of *Sphagnum* mosses or humus in forest soils with increased N deposition [Malmer, 1990; Berg and Matzner, 1997; Vitt *et al.*, 2003]. Significant changes in the microbial ecology of the peat profile with increasing N

[Gilbert *et al.*, 1998] may lead to reduced heterotrophic respiration in the upper most 5 cm of the peat profile [Williams and Silcock, 1997]. The atmospheric wet N deposition in eastern Canada (0.3–0.8 g N m⁻² yr⁻¹) is similar to that in Sweden [Aerts *et al.*, 1992] but moderate compared to that in central and western Europe with average levels of 3–4 g N m⁻² yr⁻¹ and up to 8 g N m⁻² yr⁻¹ [Houdijk and Roelofs, 1991]. Dry N deposition in eastern Canada is unknown. In areas with high levels of N deposition, many species typical of bogs and poor fens are decreasing in abundance [Greven, 1992; Tyler and Olsson, 1997]. A nutrient imbalance or a toxic effect might occur in *Sphagnum* growing under high rates of N addition, reflected for instance by increased accumulation of amino acids and iron concentrations [Ferguson *et al.*, 1984; Baxter *et al.*, 1992] or decreased photosynthesis and soluble sugar content [van der Heijden *et al.*, 2000]. Vitt *et al.* [2003] suggest that the negative impacts of net primary production on *S. fuscum* begin to occur, on average, as N deposition reaches 1.5–1.6 g N m⁻² yr⁻¹. The N-deposition values of our study sites are far below the suggested critical N-deposition values. Jauhiainen *et al.* [1994] reported that the optimal N loading for length growth of *Sphagnum fuscum* was 1 g N m⁻² yr⁻¹, and 3 g N m⁻² yr⁻¹ for biomass production. This suggests that N loading in our high N deposition region (~0.8 g N m⁻² yr⁻¹ plus dry deposition) are close to the optimal N concentrations to stimulate *Sphagnum* length growth and biomass production. Vitt *et al.* [2003] found an increase in *S. fuscum* production with increasing N deposition in Alberta, Canada. However, the production of *S. fuscum* in continental peatlands may partially be limited by water levels and not nutrients [Thormann and Bayley, 1997; Vitt *et al.*, 2003]. In our study, the aboveground vegetation biomass pattern across the sites did not indicate that the higher rates of recent C accumulation could be the result of higher biomass production. The most probable explanation for the higher C accumulation in peat rates may be

the slower decomposition rate of the organic material [Berg and Matzner, 1997].

[30] The 50-, 100-, and 150-year RERCA of hollows in eastern Canada increased consistently toward the class with the highest growing degree-days above +5°C and the highest N deposition class, but the elevated N deposition has been apparent only for the last 50 years. Therefore the higher remaining C mass in deeper hollow peat layers has to be due to something other than elevated N, that affects higher primary production and/or low decay rates throughout the peat deposits. The C loss pattern supports the conclusion that both the climate characteristics and the elevated N control the recent 50-year C accumulation in hollows. In hummocks, the significantly higher RERCA was evident only over the last 50 years. The differences in remaining C mass of deeper hummock layers were insignificant (Figure 4). The results of our study suggest that the moderately high N deposition has partly increased the present-day C accumulation both in hollows and hummocks of ombrotrophic peatlands in eastern Canada.

[31] Our results also indicated an inverse relationship between 50-year RERCA and degree-days above +5°C for hollows and hummocks (Figure 3). This result may be due to functional difference between microforms and *Sphagnum* species on ombrotrophic bogs [e.g., Malmer et al., 1994]. The primary production and decay rates of different microforms response to the temperature and the water table fluctuations. The drought and lowered water tables can cause increased oxidation and reduced primary production, resulting in a heavy C loss [Waddington and Roulet, 1996; Carroll and Crill, 1997], especially in hummocks [Alm et al., 1999].

[32] The recent (RERCA) rates were strikingly higher than the long-term (LORCA) rates of C accumulation. The LORCA average of 19 g C m⁻² yr⁻¹ is similar to many other recently published rates (17–21 g C m⁻² yr⁻¹) [Clymo et al., 1998; Vitt et al., 2000; Turunen et al., 2001, 2002] (Figure 5). Decreased LORCA with increased fire frequency shows that peatland fires (such as at sites 5 and 13) can slow peat accumulation, as individual fires can create C losses of up to 1.5 to 4 kg m⁻² [Kuhry, 1994; Pitkänen et al., 1999; Robinson and Moore, 2000]. The results of this study revealed no significant relationships between long-term C accumulation and individual climate characteristics. Also, there was only a weak relationship between LORCA and RERCA. The results indicate that the long-term C accumulation of eastern Canada has depended on a whole range of climatic, edaphic, and autogenic factors (temperature, precipitation, evapotranspiration, topography, substrate and surficial geology, groundwater flow and chemistry, peat fires). Discrimination among individual factors is difficult.

4.3. Nitrogen Accumulation

[33] The average RERNA over the past 50 years ranged from 1.4 to 3.2 g N m⁻² yr⁻¹, depending on the microsite and the N deposition area (Figure 6). These values are of the same magnitude as the N accumulation rate (2.0 g N m⁻² yr⁻¹) measured in surface bog peat in southern Sweden [Malmer and Holm, 1984], with similar wet N deposition rates to

eastern Canada [Aerts et al., 1992]. We found that the N accumulation of peat seemed to increase during the last 50 years compared to the average values for the last 150-year period (1.1–2.5 g N m⁻² yr⁻¹). Also, when comparing the remaining N mass of hummocks and hollows from 100 to 150 years, significant differences were not found between the high and low N deposition classes.

[34] Woodin and Lee [1987] and Aerts et al. [1992] showed that at high rates of N deposition the *Sphagnum* is unable to retain the entire N deposited on it, and a substantial amount of N leaches through the surface peat layers. This surplus N may be taken up by vascular plants or by microorganisms and be a partial cause of the larger vascular plant leaf biomass we found in the highest N deposition region and/or decomposition and mineralization rates of the peat [e.g., Malmer, 1990]. Our data showed that the N concentrations in deeper peat layers increased consistently toward the highest N deposition class. However, it is normal that the highest N concentrations are found in samples with more decomposed peat and thus higher dry bulk density [Malmer and Holm, 1984], and therefore it is difficult to distinguish the possible increase of N in deeper peat layers because of leaching.

[35] In summary, we have determined rates of C accumulation over the past 50 to 150 years in ombrotrophic bogs across eastern Canada, and compare them to the millennial rates. We have shown that the recent rates are correlated with environmental properties, such as climate and atmospheric N deposition between 1990 and 1996. We suggest that increased N deposition leads to larger rates of C and N accumulation in the bogs, as has been found in European forests [Kauppi et al., 1992; Berg and Matzner, 1997], and could account for some of the missing C sink in the global C budget. It is possible that fertilization responses, in particular to N, play a significant role in moderate and high anthropogenic deposition areas of North American and European peatlands. In the long term, both the positive and negative impacts for net primary production of *Sphagnum* species depend on the critical N-deposition loads as suggested by Vitt et al. [2003]. However, there are uncertainties associated with covarying properties. These uncertainties include the influence of climate, such as precipitation, temperature, and pollutant emissions, on rates of plant production and decomposition, disturbances such as fire, and pre-1990 rates of atmospheric N deposition.

[36] **Acknowledgments.** We thank Climate Change Action Fund (CCAF) and NSERC for their financial support of this research, and C. Blodau of the University of Bayreuth for the C and N analyses. J. T. received partial funding from the FCAR-supported Centre for Climate and Global Change Research for his postdoctoral fellowship.

References

- Aerts, R., B. Wallen, and N. Malmer (1992), Growth-limiting nutrients in *Sphagnum*-dominated bogs subject to low and high atmospheric nitrogen supply, *J. Ecol.*, *80*, 131–140.
- Alm, J., L. Schulman, J. Walden, H. Nykänen, P. J. Martikainen, and J. Silvola (1999), Carbon balance of a boreal bog during a year with an exceptionally dry summer, *Ecology*, *80*, 161–174.
- Appleby, P. G., and F. R. Oldfield (1978), The calculation of ²¹⁰Pb dates assuming a constant rate of supply of unsupported ²¹⁰Pb to the sediment, *Catena*, *5*, 1–8.

- Appleby, P. G., W. Shotyk, and A. Fankhauser (1997), Lead-210 age dating of three peat cores in the Jura Mountains, Switzerland, *Water Air Soil Pollut.*, *100*, 223–231.
- Baxter, R., M. J. Emes, and J. A. Lee (1992), Effects of an experimentally applied increase in ammonium on growth and amino-acid metabolism of *Sphagnum cuspidatum* Ehrh. ex. Hoffm. from differently polluted areas, *New Phytol.*, *120*, 265–274.
- Belyea, L. R. (1996), Separating the effects of litter quality and microenvironment on decomposition rates in a patterned peatland, *Oikos*, *77*, 529–539.
- Belyea, L. R., and B. G. Warner (1994), Dating of the near-surface layer of a peatland in northwestern Ontario, Canada, *Boreas*, *23*, 259–269.
- Berendse, F., et al. (2001), Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production, *Global Change Biol.*, *7*, 591–598.
- Berg, B., and E. Matzner (1997), Effect of N deposition on decomposition of plant litter and soil organic matter in forest systems, *Environ. Rev.*, *5*, 1–25.
- Botch, M. S., K. I. Kobak, T. S. Vinson, and T. P. Kolchugina (1995), Carbon pools and accumulation in peatlands of the Former Soviet Union, *Global Biogeochem. Cycles*, *9*, 37–46.
- Carroll, P. C., and P. Crill (1997), Carbon balance of a temperate poor fen, *Global Biogeochem. Cycles*, *11*, 349–356.
- Chapman, R. R., and H. F. Hemond (1982), Dinitrogen fixation by surface peat and *Sphagnum* in an ombrotrophic bog, *Can. J. Bot.*, *60*, 538–543.
- Clymo, R. S. (1984), The limits to peat bog growth, *Philos. Trans. R. Soc. London, Ser. B*, *303*, 605–654.
- Clymo, R. S., J. Turunen, and K. Tolonen (1998), Carbon accumulation in peatland, *Oikos*, *81*, 368–388.
- Damman, A. W. H. (1978), Distribution and movement of elements in ombrotrophic peat bogs, *Oikos*, *30*, 480–495.
- Damman, A. W. H. (1988), Regulation of nitrogen removal and retention in *Sphagnum* bogs and other peatlands, *Oikos*, *51*, 291–305.
- Damman, A. W. H., and J. J. Dowhan (1981), Vegetation and habitat conditions in Western Head Bog, a southern Nova Scotian plateau bog, *Can. J. Bot.*, *59*, 1343–1359.
- El-Daoushy, F., K. Tolonen, and R. Rosenberg (1982), Lead 210 and moss-increment dating of two Finnish *Sphagnum* hummocks, *Nature*, *296*, 429–431.
- Ferguson, P., R. N. Robinson, M. C. Press, and J. A. Lee (1984), Element concentrations in five *Sphagnum* species in relation to atmospheric pollution, *J. Bryol.*, *13*, 107–114.
- Gilbert, D., C. Amblard, G. Bourdier, and A.-J. Francez (1998), Short-term effect of nitrogen enrichment on the microbial communities of a peatland, *Hydrobiology*, *373/374*, 111–119.
- Glaser, P. H., and J. A. Janssens (1986), Raised bogs in eastern North America: Transitions in landforms and gross stratigraphy, *Can. J. Bot.*, *64*, 395–415.
- Gorham, E. (1990), Biotic Impoverishment in Northern Peatlands, in *The Earth in Transition: Patterns and Processes of Biotic Impoverishment*, edited by G. M. Woodwell, pp. 65–98, Cambridge Univ. Press, New York.
- Gorham, E. (1991), Northern Peatlands: Role in the carbon cycle and probable responses to climatic warming, *Ecol. Appl.*, *1*, 182–195.
- Gorham, E., S. J. Eisenreich, J. Ford, and M. V. Santelmann (1985), The chemistry of bog waters, in *Chemical Processes in Lakes*, edited by W. Stumm, pp. 339–363, Wiley-Intersci., John Wiley, New York.
- Greven, H. C. (1992), Changes in the moss flora of the Netherlands, *Biol. Conserv.*, *59*, 133–137.
- Houdijk, A. L. F. M., and J. G. M. Roelofs (1991), Deposition of acidifying and eutrophating substances in Dutch forests, *Acta Bot. Neerl.*, *40*, 245–255.
- Houghton, J. T., G. J. Jenkins, and J. J. Ephraums (1990), *Climate Change, The IPCC Scientific Assessment*, Cambridge Univ. Press, New York.
- Houghton, R. A. (1993), Is carbon accumulating in the northern temperate zone?, *Global Biogeochem. Cycles*, *7*, 611–617.
- Jauhainen, J., H. Vasander, and J. Silvola (1994), Response of *Sphagnum fuscum* to N deposition and increased CO₂, *J. Bryol.*, *18*, 83–95.
- Kauppi, P. E., K. Mielikäinen, and K. Kuusela (1992), Biomass and carbon budget of European forests, *Science*, *256*, 70–74.
- Kuhry, P. (1994), The role of fire in the development of *Sphagnum*-dominated peatlands in western Boreal Canada, *J. Ecol.*, *82*, 899–910.
- Lafleur, P. M., N. T. Roulet, and S. W. Admiral (2001), Annual cycle of CO₂ exchange at a bog peatland, *J. Geophys. Res.*, *106*, 3071–3081.
- Lindholm, T., and H. Vasander (1990), Production of eight species of *Sphagnum* at Suurisuo mire, southern Finland, *Ann. Bot. Fenn.*, *27*, 145–157.
- Livett, E. A., J. A. Lee, and J. H. Tallis (1979), Lead, zinc, and copper analysis of British blanket peats, *J. Ecol.*, *67*, 865–891.
- Malmer, N. (1990), Constant or increasing nitrogen concentrations in *Sphagnum* mosses on mires in Southern Sweden during the last few decades, *Aquilo Ser. Bot.*, *28*, 57–65.
- Malmer, N., and E. Holm (1984), Variation in the C/N-quotient of peat in relation to decomposition rate and age determination with ²¹⁰Pb, *Oikos*, *43*, 171–182.
- Malmer, N., B. M. Svensson, and B. Wallen (1994), Interactions between *Sphagnum* and field layer vascular plants in the development of peat-forming systems, *Folia Geobot. Phytotaxon.*, *29*, 483–496.
- Mitchell, P. I., W. R. Schell, A. McGarry, T. P. Ryan, J. A. Sanchez-Cabeza, and A. Vidal-Quadras (1992), Studies of the vertical distribution of ¹³⁴Cs, ¹³⁷Cs, ²³⁸Pu, ^{239,240}Pu, ²⁴¹Pu, ²⁴¹Am and ²¹⁰Pb in ombrogenous mires at mid-latitudes, *J. Radioanal. Nucl. Chem.*, *156*, 361–387.
- Moore, T. R., J. L. Bubier, S. E. Frolking, P. M. Lafleur, and N. T. Roulet (2002), Plant biomass and production and CO₂ exchange in an ombrotrophic bog, *J. Ecol.*, *90*, 25–36.
- Ohlson, M. O., and B. Dahlberg (1991), Rate of peat increment in hummocks and lawn communities on Swedish mires during the last 150 years, *Oikos*, *61*, 369–378.
- Økland, R. H., and M. Ohlson (1998), Age-depth relationships in Scandinavian surface peat: A quantitative analysis, *Oikos*, *82*, 29–36.
- Oldfield, F., P. Appleby, R. Cambray, J. Eakins, K. Barber, R. Batterby, G. Pearson, and J. Williams (1979), Pb-210, Cs-137, and Pu-239 profiles in ombrotrophic peat, *Oikos*, *33*, 40–45.
- Oldfield, F., N. Richardson, and P. G. Appleby (1995), Radiometric dating (²¹⁰Pb, ¹³⁷Cs, ²⁴¹Am) of recent ombrotrophic peat accumulation and evidence for changes in mass balance, *Holocene*, *5*, 141–148.
- Pakarinen, P., K. Tolonen, S. Heikkinen, and A. Nurmi (1983), Accumulation of metals in Finnish raised bogs, *Environ. Biogeochem.*, *35*, 377–382.
- Pitkänen, A., J. Turunen, and K. Tolonen (1999), The role of fire in the carbon dynamics of a mire, Eastern Finland, *Holocene*, *9*, 453–462.
- Post, W. M., W. R. Emanuel, P. J. Zinke, and A. G. Stangenberger (1982), Soil carbon pools and world life zones, *Nature*, *298*, 156–159.
- Robinson, S. D., and T. R. Moore (2000), The influence of permafrost and fire upon carbon accumulation in high boreal peatlands, Northwest Territories, Canada, *Arct. Alp. Res.*, *32*, 155–166.
- Shotyk, W., A. K. Cheburkin, P. G. Appleby, A. Fankhauser, and J. D. Kramers (1997), Lead in three bog profiles, Jura Mountains, Switzerland: Enrichment factors, isotopic composition, and chronology of atmospheric deposition, *Water Air Soil Pollut.*, *100*, 297–310.
- Sjörs, H. (1959), Bogs and fens in the Hudson Bay lowlands, *J. Arct. Inst. North Am.*, *12*, 1–19.
- Stuiver, M., and P. J. Reimer (1993), Extended ¹⁴C data base and revised CALIB 3.0 ¹⁴C age calibration program, *Radiocarbon*, *35*, 215–230.
- Tabatabai, M. A., and J. M. Bremer (1990), Automated instruments for determination of total carbon, nitrogen, and sulfur in soils by combustion techniques, in *Soil Analysis: Modern Instrumental Techniques*, 2nd ed., edited by K. A. Smith, pp. 261–286, Marcel Dekker, New York.
- Tans, P. P., I. Y. Fung, and T. Takahashi (1990), Observational constraints on the global atmospheric CO₂ budget, *Science*, *247*, 1431–1438.
- Thormann, M. E., and S. E. Bayley (1997), Response of above ground net primary plant production in nitrogen and phosphorous fertilization in peatlands in southern boreal, Alberta, Canada, *Wetlands*, *17*, 502–512.
- Tolonen, K., and J. Turunen (1996), Accumulation rates of carbon in mires in Finland and implications for climate change, *Holocene*, *6*, 171–178.
- Tolonen, K., R. B. Davis, and L. Widoff (1988), Peat accumulation rates in selected Maine peat deposits, Maine Geological Survey, *Dep. Conserv. Bull.*, *33*, 1–99.
- Turetsky, M. R., R. K. Wieder, C. J. Williams, and D. H. Vitt (2000), Organic matter accumulation, peat chemistry, and permafrost melting in peatlands of boreal Alberta, *Ecoscience*, *7*, 379–392.
- Turunen, J., A. Pitkänen, T. Tahvanainen, and K. Tolonen (2001), Carbon accumulation in West Siberian mires, Russia, *Global Biogeochem. Cycles*, *15*, 285–296.
- Turunen, J., E. Tomppo, K. Tolonen, and A. Reinikainen (2002), Estimating carbon accumulation rates of undrained mires in Finland: Application to boreal and subarctic regions, *Holocene*, *12*, 69–80.
- Tyler, G., and K.-A. Olsson (1997), Förändringar i Skånes flora under perioden 1938–1996—Statistisk analys av resultat från två inventeringar (Changes in the flora of Scania during the years 1938–1996), *Sven. Bot. Tidskr.*, *91*, 143–185.
- Urban, N. R., and S. J. Eisenreich (1988), Nitrogen cycling in a forested Minnesota bog, *Can. J. Bot.*, *66*, 435–449.
- Urban, N. R., S. J. Eisenreich, D. F. Grigal, and K. T. Schurr (1990), Mobility and diagenesis of Pb and ²¹⁰Pb in peat, *Geochim. Cosmochim. Acta*, *54*, 3329–3346.

- van der Heijden, E., S. K. Verbeek, and P. J. C. Kuiper (2000), Elevated atmospheric CO₂ and increasing nitrogen deposition: Effects on C and N metabolism and growth of the peat moss *Sphagnum recurvum* P. Beauv. var. mucronatum (Russ.) Warnst, *Global Change Biol.*, *6*, 201–212.
- Vet, R., C. U. Ro, and D. Ord (1999), SOE technical supplement, *Bull.* 99-3, Natl. Atmos. Chem. Database and Anal. Facil., Atmos. Environ. Serv., Environ. Can., Ontario, Canada.
- Vile, M. A., M. J. V. Novak, E. Brizova, R. K. Wieder, and W. R. Schell (1995), Historical rates of atmospheric Pb deposition using ²¹⁰Pb dated peat cores: corroboration, computation, and interpretation, *Water Air Soil Pollut.*, *79*, 89–106.
- Vile, M. A., R. K. Wieder, and M. Novak (1999), Mobility of Pb in *Sphagnum*-derived peat, *Biogeochemistry*, *45*, 35–52.
- Vitt, D. H., L. A. Halsey, I. E. Bauer, and C. Campbell (2000), Spatial and temporal trends in carbon storage of peatlands of continental western Canada through the Holocene, *Can. J. Earth Sci.*, *37*, 683–693.
- Vitt, D. H., K. Wieder, L. A. Halsey, and M. Turetsky (2003), Response of *Sphagnum fuscum* to nitrogen deposition: A case study of ombrogenous peatlands in Alberta, Canada, *Bryologist*, 235–245.
- von Post, L. (1922), Sveriges geologiska undersöknings torvinventering och några av dess hittills vunna resultat, *Sven. Mosskulturforen. Tidskr.*, *1*, 1–27.
- Waddington, J. M., and N. T. Roulet (1996), Atmosphere-wetland carbon exchanges: Scale dependency of CO₂ and CH₄ exchange on the developmental topography of a peatland, *Global Biogeochem. Cycles*, *10*, 233–245.
- Waughman, G. J., and D. J. Bellamy (1980), Nitrogen fixation and the nitrogen balance in peatland ecosystems, *Ecology*, *61*, 1185–1198.
- Wieder, R. K., M. Novak, W. R. Schell, and T. Rhodes (1994), Rates of peat accumulation over the past 200 years in five *Sphagnum*-dominated peatlands in the United States, *J. Paleolimnol.*, *12*, 35–47.
- Williams, B. L., and D. J. Silcock (1997), Nutrient and microbial changes in the peat profile beneath *Sphagnum magellanicum* in response to additions of ammonium nitrate, *J. Ecol.*, *34*, 961–970.
- Woodin, S. J., and J. A. Lee (1987), The fate of some components of acidic deposition in ombrotrophic mires, *Environ. Pollut.*, *45*, 61–72.

T. R. Moore, N. T. Roulet, and J. Turunen, Department of Geography, McGill University, 805 Sherbrooke Street West, Montréal, Québec, H3A 2K6, Canada. (tim.moore@mcgill.ca; nigel.roulet@mcgill.ca; jukka.turunen@gtk.fi)

P. J. H. Richard, Département de Géographie, Université de Montréal, C.P. 6128 Centre-ville, Montréal, Québec, H3C 3J7, Canada. (pierre.richard@umontreal.ca)